

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
10 March 2005 (10.03.2005)

PCT

(10) International Publication Number  
WO 2005/021539 A1

(51) International Patent Classification<sup>7</sup>: C07D 405/14,  
413/14, 401/12, 405/06, A61K 31/4184, A61P 1/04

[JP/JP]: Pfizer Global Research and Development, 2, Aza  
5-gochi, Taketocho-cho, Chita-gun, Aichi-ken 470-2393  
(JP).

(21) International Application Number:

PCT/IB2004/002741

(74) Agents: FULLER, Grover, F., Jr. et al.; Pfizer Inc., P.O.  
Box 1027, St. Louis, MO 63006 (US).

(22) International Filing Date: 20 August 2004 (20.08.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/500,144 3 September 2003 (03.09.2003) US

(71) Applicant (for JP only): PFIZER JAPAN, INC. [JP/JP];  
Shinjuku Bunka Quint Bldg., 3-22-7, Yoyogi, Shibuya-ku,  
Tokyo 151-8589 (JP).

(71) Applicant (for all designated States except JP, US):  
PFIZER INC. [US/US]; 235 East 42nd Street, New York,  
NY 10017 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): IGUCHI, Satoru  
[JP/JP]; Pfizer Global Research and Development, 2, Aza  
5-gochi, Taketocho-cho, Chita-gun, Aichi-ken 470-2393  
(JP). KATSU, Yasuhiro [JP/JP]; Pfizer Global Re-  
search and Development, 2, Aza 5-gochi, Taketocho-cho,  
Chita-gun, Aichi-ken 470-2393 (JP). SONE, Hiroki  
[JP/JP]; Pfizer Global Research and Development, 2, Aza  
5-gochi, Taketocho-cho, Chita-gun, Aichi-ken 470-2393  
(JP). UCHIDA, Chikara [JP/JP]; Pfizer Global Re-  
search and Development, 2, Aza 5-gochi, Taketocho-cho,  
Chita-gun, Aichi-ken 470-2393 (JP). KOJIMA, Takashi

(81) Designated States (unless otherwise indicated, for every  
kind of national protection available): AE, AG, AL, AM,  
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,  
CO, CR, CU, CZ, DE, DK, DM, DO, EC, EE, EG, ES, FI,  
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,  
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,  
MG, MK, MN, MW, MX, MZ, NA, NL, NO, NZ, OM, PG,  
PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM,  
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM,  
ZW.


(84) Designated States (unless otherwise indicated, for every  
kind of regional protection available): ARIPO (BW, GH,  
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,  
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,  
FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI,  
SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,  
GW, ML, MR, NE, SN, TD, TG).

#### Published:

- with international search report
- before the expiration of the time limit for amending the  
claims and to be republished in the event of receipt of  
amendments

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

(54) Title: BENZIMIDAZOLONE COMPOUNDS HAVING 5-HT<sub>4</sub> RECEPTOR AGONISTIC ACTIVITY



(I)

(57) Abstract: This invention provides a compound of the formula (I) or a pharmaceutically acceptable salt thereof, and compositions containing such compounds and the use of such compounds for the manufacture of medicament for gastroesophageal reflux disease, gastrointestinal disease, gastric motility disorder, non-ulcer dyspepsia, functional dyspepsia, irritable bowel syndrome (IBS), constipation, dyspepsia, esophagitis, gastroesophageal disease, nausea, central nervous system disease, Alzheimer's disease, cognitive disorder, emesis, migraine,

neurological disease, pain, cardiovascular disorders, cardiac failure, heart arrhythmia, diabetes and apnea syndrome. These compounds have 5-HT<sub>4</sub> receptor agonistic activity, and thus are useful for the treatment of gastroesophageal reflux disease, non-ulcer dyspepsia, functional dyspepsia, irritable bowel syndrome or the like in mammalian, especially humans.

**BENZIMIDAZOLONE COMPOUNDS**  
**HAVING 5-HT<sub>4</sub> RECEPTOR AGONISTIC ACTIVITY**

**Technical Field**


This invention relates to novel benzimidazolone compounds. These  
5 compounds have selective 5-HT<sub>4</sub> receptor agonistic activity. The present invention  
also relates to a pharmaceutical composition, a method of treatment and use  
comprising the above compounds for the treatment of disease conditions mediated by  
5-HT<sub>4</sub> receptor activity.

**Background Art**

10 In general, 5-HT<sub>4</sub> receptor agonists are found to be useful for the treatment of  
a variety of diseases such as gastroesophageal reflux disease, gastrointestinal disease,  
gastric motility disorder, non-ulcer dyspepsia, functional dyspepsia, irritable bowel  
syndrome (IBS), constipation, dyspepsia, esophagitis, gastroesophageal disease,  
nausea, central nervous system disease, Alzheimer's disease, cognitive disorder,  
15 emesis, migraine, neurological disease, pain, cardiovascular disorders, cardiac failure,  
heart arrhythmia, diabetes and apnea syndrome (See *TiPs*, 1992, 13, 141; Ford A. P.  
D. W. et al., *Med. Res. Rev.*, 1993, 13, 633; Gullikson G. W. et al., *Drug Dev.*  
*Res.*, 1992, 26, 405; Richard M. Eglén et al., *TiPS*, 1995, 16, 391; Bockaert J. Et al.,  
*CNS Drugs*, 1, 6; Romanelli M. N. et al., *Arzheim Forsch./Drug Res.*, 1993, 43,  
20 913; Kaumann A. et al., *Naunyn-Schmiedeberg's*. 1991, 344, 150; and Romanelli M.  
N. et al., *Arzheim Forsch. /Drug Res.*, 1993, 43, 913). Also, Mosapride is known to be  
useful for the treatment of diabetes.


It would be desirable if there were provided 5-HT<sub>4</sub> receptor agonists which  
have more 5HT<sub>4</sub> receptor agonistic activities.

25 US5223511 discloses benzimidazole compounds as 5-HT<sub>4</sub> receptor  
antagonists. Especially, compounds represented by the following formula is  
disclosed:




**Compound A**

WO93/18027 discloses benzimidazolone compounds as 5-HT<sub>4</sub> receptor antagonists. Especially, compounds represented by the following formula is disclosed:




**Compound B**

5 WO99/17772 discloses benzimidazolone compounds as 5-HT<sub>4</sub> receptor agonists and/or antagonists. Especially, compounds represented by the following formula is disclosed:



**Compound C**

10 WO94/00449 discloses benzimidazolone compounds as 5-HT<sub>4</sub> agonists or antagonists and/or 5-HT<sub>3</sub> antagonists. Especially, compounds represented by the following formula is disclosed:



**Compound D**

There is a need to provide new 5-HT<sub>4</sub> agonists that are good drug candidates. In particular, preferred compounds should bind potently to the 5-HT<sub>4</sub> receptor whilst showing little affinity for other receptors and show functional activity as agonists. They should be well absorbed from the gastrointestinal tract, be metabolically stable and possess favorable pharmacokinetic properties. When targeted against receptors in the central nervous system they should cross the blood brain barrier freely and when targeted selectively against receptors in the peripheral nervous system they should not cross the blood brain barrier. They should be non-toxic and demonstrate few side-

15

20

effects. Furthermore, the ideal drug candidate will exist in a physical form that is stable, non-hygroscopic and easily formulated.


### **Brief Disclosure of the Invention**

5           It has now surprisingly been found that compounds of this invention have strong selective 5-HT<sub>4</sub> agonistic activity, and thus are useful for the treatment of disease conditions mediated by 5-HT<sub>4</sub> activity such as gastroesophageal reflux disease, gastrointestinal disease, gastric motility disorder, non-ulcer dyspepsia, functional dyspepsia, irritable bowel syndrome (IBS), constipation, dyspepsia, esophagitis, 10 gastroesophageal disease, nausea, central nervous system disease, Alzheimer's disease, cognitive disorder, emesis, migraine, neurological disease, pain, cardiovascular disorders, cardiac failure, heart arrhythmia, diabetes and apnea syndrome (especially caused by an opioid administration).

          Further, the compounds of the present invention show a reduced QT 15 prolongation by introducing a polar group into R<sup>3</sup> of the formula (I). QT prolongation is known to have a potential liability to produce fatal cardiac arrhythmias of Torsades de Pointes (TdP). The ability to prolong the cardiac action potential duration was identified as being due to an action at the HERG potassium channel. For example, drugs withdrawn from the market due to QT prolongation, such as Cisapride and 20 Terfenadine, are known to be potent HERG potassium channel blocker (Expert Opinion of Pharmacotherapy.; 2, pp947-973, 2000). Inhibitory activity at HERG channel was estimated from affinity for HERG type potassium channel was investigated by checking [<sup>3</sup>H]dofetilide binding, which can predict inhibitory activity at HERG channel (Eur. J. Pharmacol., 430, pp147-148, 2001).

25           The compounds of the present invention may show less toxicity, good absorption, distribution, good solubility, low protein binding affinity, less drug-drug interaction, and good metabolic stability.

          The present invention provides a compound of the following formula (I) or a pharmaceutically acceptable salt thereof.



wherein

**Het** represents a heterocyclic group having one nitrogen atom, to which B binds directly, and from 4 to 7 carbon atoms, and said heterocyclic group being

- 5 unsubstituted or substituted by 1 to 4 substituents independently selected from the group consisting of substituents  $\alpha^1$ ;

**A** represents an alkylene group having from 1 to 4 carbon atoms;

**B** represents a covalent bond or an alkylene group having from 1 to 5 carbon atoms, and said alkylene group being unsubstituted or substituted by an oxo group when  $R^3$

- 10 represents a heterocyclic group;

$R^1$  represents an isopropyl group or a cyclopentyl group;

$R^2$  independently represents a halogen atom or an alkyl group having from 1 to 4 carbon atoms;  $m$  is 0, 1, 2, 3 or 4; and

$R^3$  represents

- 15 (i) a cycloalkyl group having from 3 to 8 carbon atoms, and said cycloalkyl group being substituted by 1 to 5 substituents independently selected from the group consisting of substituents  $\alpha^2$ , or
- (ii) a heterocyclic group having from 3 to 8 atoms, and said heterocyclic group being unsubstituted or substituted by 1 to 5 substituents independently
- 20 selected from the group consisting of substituents  $\beta$ ,

said substituents  $\alpha^1$  are independently selected from a hydroxy group and an amino group;

said substituents  $\alpha^2$  are independently selected from a hydroxy group, an amino group, a hydroxy-substituted alkyl group having from 1 to 4 carbon atoms, a carboxyl group and an alkoxy group having from 1 to 4 carbon atoms; and

said substituents  $\beta$  are independently selected from a hydroxy group, a hydroxy-


substituted alkyl group having from 1 to 4 carbon atoms, a carboxyl group, an amino group, an alkyl group having from 1 to 4 carbon atoms, an amino-substituted alkyl group having from 1 to 4 carbon atoms and a carbamoyl group,

Also, the present invention provides a pharmaceutical composition for the  
5 treatment of disease conditions mediated by 5-HT<sub>4</sub> receptor, in a mammalian subject, which comprises administering to said subject a therapeutically effective amount of a compound of formula (I) or pharmaceutically acceptable salts thereof.

Further, the present invention also provides a pharmaceutical composition for the treatment of diseases selected from gastroesophageal reflux disease,  
10 gastrointestinal disease, gastric motility disorder, non-ulcer dyspepsia, functional dyspepsia, irritable bowel syndrome (IBS), constipation, dyspepsia, esophagitis, gastroesophageal disease, nausea, central nervous system disease, Alzheimer's disease, cognitive disorder, emesis, migraine, neurological disease, pain, cardiovascular disorders, cardiac failure, heart arrhythmia, diabetes and apnea  
15 syndrome, or the like, which comprises a therapeutically effective amount of the benzimidazolone compound of formula (I) or its pharmaceutically acceptable salt together with a pharmaceutically acceptable carrier.

Also, the present invention provides a method of the treatment of a mammal, including a human, to treat a disease conditions mediated by 5-HT<sub>4</sub> receptor, in a  
20 mammalian subject, which comprises administering to said subject a therapeutically effective amount of a compound of formula (I) or pharmaceutically acceptable salts thereof. Further, the present invention provides a method for the treatment of the disease conditions as mentioned above. Furthermore, the present invention provides use of the compound of formula (I) or pharmaceutically acceptable salts thereof in the  
25 manufacture of a medicament for the treatment of disease conditions mediated by 5-HT<sub>4</sub> receptor activity, in a mammalian subject. The conditions mediated by 5-HT<sub>4</sub> receptor activity include those diseases or disorders described as above.

Also, the present invention provides a compound of the following formula (2-A') or a salt thereof:



wherein

**R<sup>a</sup>** represents a hydrogen atom or a N-protecting group;

**Het** represents a heterocyclic group having one nitrogen atom, to which B binds directly, and from 4 to 7 carbon atoms, and said heterocyclic group being unsubstituted or substituted by 1 to 4 substituents independently selected from the group consisting of substituents  $\alpha^1$ ;

**A** represents an alkylene group having from 1 to 4 carbon atoms;

**B** represents a covalent bond or an alkylene group having from 1 to 5 carbon atoms, and said alkylene group being unsubstituted or substituted by an oxo group when **R<sup>3</sup>** represents a heterocyclic group;

**R<sup>3</sup>** represents

- (i) a cycloalkyl group having from 3 to 8 carbon atoms, and said cycloalkyl group being substituted by 1 to 5 substituents independently selected from the group consisting of substituents  $\alpha^2$ , or
- (ii) a heterocyclic group having from 3 to 8 atoms, and said heterocyclic group being unsubstituted or substituted by 1 to 5 substituents independently selected from the group consisting of substituents  $\beta$ ,


**said substituents  $\alpha^1$**  are independently selected from a hydroxy group and an amino group;

**said substituents  $\alpha^2$**  are independently selected from a hydroxy group, an amino group, a hydroxy-substituted alkyl group having from 1 to 4 carbon atoms, a carboxyl group and an alkoxy group having from 1 to 4 carbon atoms; and

**said substituents  $\beta$**  are independently selected from a hydroxy group, a hydroxy-substituted alkyl group having from 1 to 4 carbon atoms, a carboxyl group, an amino group, an alkyl group having from 1 to 4 carbon atoms, an amino-substituted alkyl group having from 1 to 4 carbon atoms and a carbamoyl group,

#### Detailed Description of the Invention

As used herein, the term "**heterocyclic**" of "Het" means a heterocyclic group having one nitrogen atom and from 4 to 7 carbon atoms such as



As used herein, the term "**alkylene**" in "A" means straight or branched chain saturated radicals having 1 to 4 carbon atoms, including, but not limited to methylene, ethylene, *n*-propylene, isopropylene, *n*-butylene, isobutylene, *sec*-butylene, *tert*-butylene. The "**alkylene**" in "A" represents preferably a methylene group, an ethylene group or a propylene group; more preferably a methylene group or an ethylene group; most preferably a methylene group.

10

As used herein, the term "**alkylene**" in "B" means straight or branched chain saturated radicals having 1 to 5 carbon atoms, including, but not limited to methylene, ethylene, *n*-propylene, isopropylene, *n*-butylene, isobutylene, *sec*-butylene, *tert*-butylene, *n*-pentylene, isopentylene, *sec*-pentylene, *tert*-pentylene. The "**alkylene**" in "B" represents preferably an alkylene group having from 1 to 4 carbon atoms; more preferably an alkylene group having from 1 to 3 carbon atoms; much more preferably a methylene group or an ethylene group; further more preferably a methylene group.

15

As used herein, the term "**halogen**" in "R<sup>2</sup>" means fluoro, chloro, bromo and iodo, preferably fluoro or chloro.

20

As used herein, the term "**alkyl**" in "R<sup>2n</sup>", "**alkyl**" of "a hydroxy-substituted alkyl group" and "an alkoxy group having from 1 to 4 carbon atoms" in "substituents  $\alpha^{2n}$ ", "**alkyl**" in "substituents  $\beta$ "; and "**alkyl**" of "a hydroxy-substituted alkyl group" and "an amino-substituted alkyl group" in "substituents  $\beta$ " mean straight or branched chain saturated radicals having 1 to 4 carbon atoms, including, but not limited to methyl, ethyl, *n*-propyl, isopropyl, *n*-butyl, isobutyl, *sec*-butyl, *tert*-butyl.

25

As used herein, the term "**cycloalkyl**" in "R<sup>3</sup>" means cyclic alkyl group having 3 to 8 carbon atoms, such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl,




cycloheptyl, cyclooctyl, and etc.

As used herein, the term **“heterocyclic”** of “R<sup>3</sup>” means a heterocyclic ring which has one or more hetero atoms in the ring, preferably has 2 to 6 carbon atoms and 1 to 3 heteroatoms, including aziridinyl, azetidiny, piperidinyl,

- 5 morpholinyl(including morpholino), pyrrolidinyl, pyrazolidinyl, piperazinyl, tetrahydropyrazolyl, pyrazolinyl, tetrahydropyranyl and etc.


The term **“treating”**, as used herein, refers to reversing, alleviating, inhibiting the progress of, or preventing the disorder or condition to which such term applies, or one or more symptoms of such disorder or condition. The term “treatment” as used  
10 herein refers to the act of treating, as “treating” is defined immediately above.

A preferred compound of formula (I) of this invention is that wherein  
**Het** represents a heterocyclic group selected from



- said heterocyclic group being unsubstituted or substituted by 1 to 3 substituents  
15 independently selected from the group consisting of substituents  $\alpha^1$ ; and  
**A** represents an alkylene group having from 1 to 3 carbon atoms.

A more preferred compound of formula (I) of this invention is that wherein




- Het** represents a group of formula  
and this group being unsubstituted or substituted by one substituent selected from the  
20 group consisting of substituents  $\alpha^1$ ;  
**A** represents an alkylene group having from 1 to 2 carbon atoms;  
**B** represents an alkylene group having from 1 to 4 carbon atoms, and said alkylene group being unsubstituted or substituted by an oxo group when R<sup>3</sup> represents a heterocyclic group;  
25 **R<sup>2</sup>** independently represents a halogen atom or an alkyl group having from 1 to 2 carbon atoms; **m** is 0, 1 or 2; and

$R^3$  represents

- (i) a cycloalkyl group having from 4 to 7 carbon atoms, and said cycloalkyl group being substituted by 1 to 3 substituents independently selected from the group consisting of substituents  $\alpha^2$ , or
- 5 (ii) a heterocyclic group having from 4 to 7 atoms, and said heterocyclic group being unsubstituted or substituted by 1 to 3 substituents independently selected from the group consisting of substituents  $\beta$ .

Also, a further preferred compound of formula (I) of this invention is the compound or its pharmaceutically acceptable salt wherein



10 **Het** represents a group of formula

and this group being unsubstituted or substituted by one substituent selected from the group consisting of substituents  $\alpha^1$ ;

**A** represents a methylene group;

**B** represents an alkylene group having from 1 to 2 carbon atoms;

15  $R^1$  represents an isopropyl group;

$R^2$  independently represents a fluorine atom, a chlorine atom or a methyl; and

$R^3$  represents

- (i) a cycloalkyl group having from 5 to 7 carbon atoms, and said cycloalkyl group being substituted by 1 to 2 substituents independently selected from the group consisting of substituents  $\alpha^2$ , or
- 20 (ii) a heterocyclic group having from 5 to 7 atoms, and said heterocyclic group being unsubstituted or substituted by 1 to 2 substituents independently selected from the group consisting of substituents  $\beta$ ,


said substituents  $\alpha^2$  are independently selected from a hydroxy group, an amino

25 group and an alkoxy group having from 1 to 2 carbon atoms; and

said substituents  $\beta$  are independently selected from a hydroxy group, a hydroxy-substituted alkyl group having from 1 to 2 carbon atoms, a carboxyl group, an amino group, an amino-substituted alkyl group having from 1 to 2 carbon atoms and a carbamoyl group.

30 A further preferred compound of formula (I) of this invention is the compound

Most preferred compounds of formula (I) of this invention is the compound or its pharmaceutically acceptable salt, wherein



**Het** represents a group of formula

**A** represents a methylene group;

5 **B** represents a methylene group;

**R<sup>1</sup>** represents an isopropyl group;

**R<sup>2</sup>** represents a fluorine atom; **m** is 0; and

**R<sup>3</sup>** represents

- (i) a cyclohexyl group substituted by 1 or 2 hydroxy group (especially dihydroxycyclohexyl), or
- 10 (ii) a tetrahydropyran group substituted by 1 or 2 hydroxy group (especially hydroxytetrahydropyranyl).

In the compounds of formula (I) or the pharmaceutically acceptable salt, **R<sup>2</sup>** preferably represents a fluorine atom, a chlorine atom, a methyl group or an ethylene group; more preferably a fluorine atom, a chlorine atom, a methyl group; most

15 preferably a fluorine atom.

In the compounds of formula (I) or the pharmaceutically acceptable salt, **m** is preferably 0, 1 or 2; more preferably 0 or 1; much more preferably 0.


Preferred individual compound of this invention is:

- 20 *N*-({1-[(4-hydroxytetrahydro-2*H*-pyran-4-yl)methyl]piperidin-4-yl}methyl)-3-isopropyl-2-oxo-2,3-dihydro-1*H*-benzimidazole-1-carboxamide;
- N*-({1-[(*trans*-1,4-dihydroxyhexyl)methyl]piperidin-4-yl}methyl)-3-isopropyl-2-oxo-2,3-dihydro-1*H*-benzimidazole-1-carboxamide;
- N*-({1-[(*cis*-1,4-dihydroxyhexyl)methyl]piperidin-4-yl}methyl)-3-isopropyl-2-oxo-
- 25 2,3-dihydro-1*H*-benzimidazole-1-carboxamide; and
- 6-fluoro-*N*-({1-[(4-hydroxytetrahydro-2*H*-pyran-4-yl)methyl]piperidin-4-yl}methyl)-3-isopropyl-2-oxo-2,3-dihydro-1*H*-benzimidazole-1-carboxamide,
- or a pharmaceutically acceptable salt thereof.

A preferred compound of formula (2-A') of this invention is that wherein

- 30 **R<sup>a</sup>** represents a hydrogen atom or a *t*-butoxycarbonyl group;

or its pharmaceutically acceptable salt wherein



**Het** represents a group of formula

**A** represents a methylene group;

**B** represents a methylene group;


5 **R<sup>1</sup>** represents an isopropyl group;

**R<sup>2</sup>** represents a fluorine atom; **m** is 0 or 1; and

**R<sup>3</sup>** represents

- (i) a cycloalkyl group having from 5 to 6 carbon atoms, and said cycloalkyl group being substituted by 1 to 2 substituents independently selected from the group consisting of substituents  $\alpha^2$ , or
- 10 (ii) a heterocyclic group having from 5 to 6 atoms, and said heterocyclic group being unsubstituted or substituted by 1 to 2 substituents independently selected from the group consisting of substituents  $\beta$ ,
- said substituents  $\alpha^2$  are independently selected from a hydroxy group and an amino group;
- 15 said substituents  $\beta$  are independently selected from a hydroxy group and an amino group.

A further preferred compound of formula (I) of this invention is the compound or its pharmaceutically acceptable salt, wherein



20 **Het** represents a group of formula

**A** represents a methylene group;


**B** represents a methylene group;

**R<sup>1</sup>** represents an isopropyl group;

**R<sup>2</sup>** represents a fluorine atom; **m** is 0; and

25 **R<sup>3</sup>** represents

- (i) a cyclohexyl group substituted by 1 to 2 substituents independently selected from a hydroxy group or an amino group, or
- (ii) a heterocyclic group having from 6 atoms, and said heterocyclic group being substituted by a hydroxy group or an amino group.



Het represents a group of formula

A represents an methylene group; B represents an methylene group; and  
 R<sup>3</sup> represents hydroxytetrahydropyranyl or dihydroxycyclohexyl.

## 5 General Synthesis

The compounds of the present invention may be prepared by a variety of processes well known for the preparation of compounds of this type, for example as shown in the following reaction Schemes. Unless otherwise indicated R<sup>1</sup> through R<sup>3</sup> and m in the following reaction Schemes and discussion are defined as above. The term "protecting group", as used hereinafter, means a hydroxy or amino protecting group which is selected from typical hydroxy or amino protecting groups described in Protective Groups in Organic Synthesis edited by T. W. Greene *et al.* (John Wiley & Sons, 1991); All starting materials in the following general syntheses may be commercially available or obtained by conventional methods known to those skilled in the art.



The compound of formula (I), wherein Het is

the following synthesis. And the compound of formula (I), wherein Het is other than



, can be prepared by a similar manner or a method known to a skilled person.


In Steps 1a, 1b, 1d, 2a, 2c, 2e, 3a, 3c, 3d of the following schemes, each reaction is preferably carried out in the presence of a base. There is no particular restriction on the nature of the bases used, and any base commonly used in reactions of this type may equally be used here. The base employed includes, for example, alkali metal hydroxides, such as lithium hydroxide, sodium hydroxide and potassium hydroxide; alkali metal carbonates such as sodium carbonate and potassium carbonate; alkali metal hydrides such as sodium hydride, potassium hydride and lithium hydride; alkali metal alkoxides such as sodium methoxide, sodium ethoxide, potassium t-butoxide and lithium methoxide; alkylolithiums such as butyllithium and

- methylolithium; lithium amides such as lithium diethylamides, lithium diisopropylamide and lithium bis(trimethylsilyl)amide; alkali metal hydrogencarbonates such as sodium hydrogencarbonate and potassium hydrogencarbonate; and tertiary organic amines such as triethylamine, dimethylaniline, pyridine, 4-dimethylaminopyridine, 1,5-diazabicyclo[4.3.0]non-5-ene, 1,8-diazabicyclo[5.4.0]undec-7-ene and N,N-diisopropylethylamine.

### Synthesis of Benzimidazolone (1-A):

The following reaction Schemes illustrate the preparation of benzimidazolone compounds of formula 1-A.

#### 10 **Scheme 1a:**



In the above formulae, Z represents 'halo', such as a chlorine, bromine or iodine atom.

#### Step 1a

- In step 1a, an amine compound of formula 1-3 can be prepared by the reductive amination of the alkanone compound (having from 1 to 4 carbon atoms) with an amine compound of formula 1-1 in the presence or absence of a reducing agent or a metal agent in an inert solvent.

The reaction is normally and preferably effected in the presence of a solvent.

- There is no particular restriction on the nature of the solvent to be employed, provided

that it has no adverse effect on the reaction or on the reagents involved and that it can dissolve the reagents, at least to some extent. Examples of suitable aqueous or non-aqueous organic solvents include: alcohols, such as methanol, ethanol or isopropanol; ethers, such as tetrahydrofuran(THF), dimethoxyethane or dioxane; acetonitrile; *N,N'*-dimethylformamide; dimethylsulfoxide; acetic acid; and halogenated hydrocarbon, such as dichloromethane, dichloroethane or chloroform.

The reaction can take place over a wide range of temperatures, and the precise reaction temperature is not critical to the invention. The preferred reaction temperature will depend upon such factors as the nature of the solvent, and the starting material or reagent used. However, in general, it is convenient to carry out the reaction with reducing agents at a temperature of from  $-78^{\circ}\text{C}$  to  $100^{\circ}\text{C}$ , more preferably from about  $-20^{\circ}\text{C}$  to  $60^{\circ}\text{C}$ . The time required for the reaction may also vary widely, depending on many factors, notably the reaction temperature and the nature of the reagents and solvent employed. However, provided that the reaction is effected under the preferred conditions outlined above, a period of 5 minutes to 1 week, more preferably 30 minutes to 24 hours, will usually suffice. In the case of the reaction with metal reagents, it is convenient to carry out the reaction at a temperature of from  $20^{\circ}\text{C}$  to  $100^{\circ}\text{C}$ , preferably from about  $20^{\circ}\text{C}$  to  $60^{\circ}\text{C}$  for 10 minutes to 48 hours, preferably 30 minutes to 24 hours.

Suitable reducing reagents are those typically used in the reduction including, for example, sodium borohydride, sodium cyanoborohydride or sodium triacetoxyborohydride.

The combination of metal reagents and hydrogen gas can be also employed as reducing reagent. Example of suitable metal reagents include palladium-carbon, palladiumhydroxide-carbon, platinumoxide, platinum-carbon, ruthenium-carbon, rhodium-aluminumoxide and tris[triphenylphosphine] rhodiumchloride. The reduction with metal reagents may be carried out under hydrogen atmosphere at a pressure ranging from 1 to 100 atm, preferably from 1 to 10 atm.

This reduction can be carried out after formation of the corresponding enamine of the alkanone compound or imine of the alkanone compound in a reaction-inert solvent such as benzene, toluene, or xylene at a temperature in the range from  $20$  to  $130^{\circ}\text{C}$  for 1 hour to 1 week.

Alternatively, the compound of formula 1-3 can be prepared by alkylation of the compound of formula 1-1 with an alkyl halide of formula of  $Z-R^1$  wherein Z is halo (halo is chloro, bromo, or iodo) as essentially the same condition as below (Step 1d), preferably in the presence of a base.

5    Step 1b

In this step, a compound of formula 1-3 can be prepared by alkylation of a compound of formula 1-2 with compound of formula  $R^1-NH_2$ .

The reaction can take place over a wide range of temperatures, and the precise reaction temperature is not critical to the invention. The preferred reaction temperature will depend upon such factors as the nature of the solvent, and the starting material or reagent used. However, in general, it is convenient to carry out the reaction at a temperature of from 0 °C to 150 °C, more preferably from 20 °C to 120 °C. The time required for the reaction may also vary widely, depending on many factors, notably the reaction temperature and the nature of the reagents and solvent employed. However, provided that the reaction may be effected under the preferred conditions outlined above, a period of from 5 minutes to 48 hours, more preferably from 30 minutes to 24 hours, will usually suffice.

15    Step 1c

A compound of formula 1-4 can be prepared by reduction of a compound of formula 1-3 with a suitable reducing agent, such as sodium borohydride ( $NaBH_4$ ), lithium aluminumhydride (LAH), diborane, hydrogen and a metal catalyst, iron and hydrochloric acid, stannic chloride and hydrochloric acid, zinc and hydrochloric acid, formic acid, borane dimethylsulfide complex, borane-THF, (preferably hydrogen and a metal catalyst), usually in excess, in a reaction inert solvent such as methanol, ethanol, propanol, butanol, tetrahydrofuran (THF) (preferably methanol or ethanol), generally at temperature of -78 °C to 60 °C, preferably from about 0 °C to 45 °C for 5 minutes to 24 hours, preferably 60 minutes to 12 hours.

25    Step 1d

In step 1d, an amine compound of formula 1-4 can be prepared by the reductive amination of the alkanone compound with an amine compound of formula 1-5 in a similar condition in step 1a.

Alternatively, a compound of formula 1-4 can be prepared by alkylation of a



compound of formula 1-5 with compound of formula  $Z-R^1$ .

The reaction can take place over a wide range of temperatures, and the precise reaction temperature is not critical to the invention. The preferred reaction temperature will depend upon such factors as the nature of the solvent, and the

- 5 starting material or reagent used. However, in general, it is convenient to carry out the reaction at a temperature of from 0 °C to 120 °C, more preferably from 0 °C to 70 °C. The time required for the reaction may also vary widely, depending on many factors, notably the reaction temperature and the nature of the reagents and solvent employed. However, provided that the reaction may be effected under the preferred conditions
- 10 outlined above, a period of from 5 minutes to 48 hours, more preferably from 30 minutes to 24 hours, will usually suffice.

#### Step 1e


A compound of formula 1-A can be prepared by cyclization of a compound of formula 1-4 with a suitable carbonylating agent such as carbonyldiimidazole,

15 trichloromethyl chloroformate, triphosgene and urea (preferably carbonyldiimidazole), usually in excess, in a reaction inert solvent such as dimethoxyethane, dioxane, acetonitrile, *N,N'*-dimethylformamide, dimethylsulfoxide, dichloromethane, dichloroethane, chloroform, or tetrahydrofuran (THF) (preferably THF), generally at temperature of -78 °C to 120 °C, preferably from about 20 °C to 100 °C for 5 minutes

20 to 24 hours, preferably 60 minutes to 12 hours.

Alternatively, the compound of 1-A (wherein  $R^1$  is isopropyl as shown in Scheme 1b) can be prepared from an alkenyl-benzimidazolone compound of formula 1-6 according to the following Scheme 1b in a reaction condition known to a skilled person.

#### 25 **Scheme 1b:**




#### Synthesis of Amine Moiety (2-A):

The following reaction Schemes illustrate the preparation of piperidine

compounds of formula (2-A).

**Scheme 2a:**




- In the above formulae, PG represents a protecting group. The term “protecting group”, as used herein, means an amino protecting group which is selected from typical amino protecting groups described in *Protective Groups in Organic Synthesis* edited by T. W. Greene *et al.* (John Wiley & Sons, 1991). Typical amino protecting groups include benzyl,  $\text{C}_2\text{H}_5\text{O}(\text{C}=\text{O})$ -,  $\text{CH}_3(\text{C}=\text{O})$ -, *tert*-butyldimethylsilyl (TBS), *tert*-butyldiphenylsilyl, benzyloxycarbonyl represented as Z and *tert*-butoxycarbonyl represented as t-Boc or Boc.

- A compound of formula 2-2 can be prepared by alkylation or reductive amination of a compound of formula 2-1 with a compound of formula alkyl- $\text{R}^3$ , halo- $\text{R}^3$ , or  $\text{H}(\text{C}=\text{O})\text{-R}^3$  in a similar condition to step 1a. When  $\text{B-R}^3$  represents 4-hydroxytetrahydropyranylmethyl, this alkylation can be done by using a 1,6-dioxaspiro[2.5]octane compound.

Then, this reaction is followed by deprotection to obtain a compound of formula 1-A. This deprotection may be carried out according to procedures known to those skilled in the art to give the compound of formula of 2-A.

- Alternatively, the compound of formula (2-A) can be prepared from a piperidine compound of formula 2-3 according to the following Scheme 2b with a reaction condition known to a skilled person.

**Scheme 2b:**




( $\text{A}^1$  is a covalent bond or  $\text{C}_{1-3}$  alkylene).

- For example, in step 2c, the compound 2-4 may be prepared by alkylation or reductive amination in essentially the same condition as one described in step 2a of Scheme 2a. Then, the reduction in step 2d may be carried out in the presence of a reducing reagent such as  $\text{LiAlH}_4$  in a reaction inert solvent such as THF. Suitable

reaction temperature ranges from about -78 °C to about 100 °C, preferably from about -30 °C to about 40 °C.

- The compound of formula (1-A) can be prepared from a piperidine compound of formula 2-5 according to the following Scheme 2c with a reaction condition known to a skilled person

**Scheme 2c:**




(A¹ is a covalent bond or C<sub>1-3</sub> alkylene)

- For example, in step 2e, the compounds 2-6 may be prepared by alkylation or reductive amination in a similar condition to one described in step 2a of scheme 2a.
- Then, the reduction in step 2f may be carried out in the presence of a H<sub>2</sub> and a hydrogenation catalyst such as PtO<sub>2</sub> in a reaction inert solvent such as THF.
- Suitable reaction temperature ranges from about -78 °C to about 100 °C, preferably from about -30 °C to about 40 °C.

**Synthesis of the compound of formula (I):**

- The following reaction Schemes illustrate the preparation of benzimidazolone compounds of formula I.

**Scheme 3a:**



**Step 3a:**

- A compound of formula 3-1 can be prepared by carbonylation of a compound of formula 1-A with a compound of formula 2-A in the presence of a suitable

carbonylating agent such as carbonyldiimidazole, trichloromethyl chloroformate, triphosgene, 4-nitrophenyl chloroformate, or urea (preferably triphosgene), usually in excess, in a reaction inert solvent such as, dimethoxyethane, dioxane, acetonitrile, *N,N'*-dimethylformamide, dimethylsulfoxide, dichloromethane, dichloroethane, tetrahydrofuran (THF), benzene, toluene, or chloroform (preferably THF), generally at temperature of -78 °C to 120 °C, preferably from about 0 °C to 90 °C for 5 minutes to 24 hours, preferably 60 minutes to 12 hours.

#### Step 3b:


A compound of formula 3-2 is prepared by deprotection of a compound of formula 3-1 with an acid such as hydrochloride,

#### Step 3c:

A compound of formula (Ia) can be prepared by alkylation or reductive amination in a similar condition to one described in step 2a of Scheme 2a.

Alternatively, the compound of formula (Ia) can be prepared from alkyl-benzimidazolone compounds according to the following Scheme 3b in a reaction condition known to a skilled person.

#### Scheme 3b:




For example, in step 3d, the compound of formula 1-A can be reacted with a compound of formula 2-A in the presence of a carbonylating agent such as carbonyldiimidazole, trichloromethyl chloroformate, triphosgene, 4-nitrophenyl chloroformate, or urea (preferably triphosgene), usually in excess, in a reaction inert solvent such as dimethoxyethane, dioxane, acetonitrile, *N,N'*-dimethylformamide, dimethylsulfoxide, dichloromethane, dichloroethane, tetrahydrofuran (THF), benzene, toluene, or chloroform (preferably THF), generally at temperature of -78 °C to 120 °C, preferably from about 0 °C to 90 °C for 5 minutes to 24 hours, preferably 60 minutes to 12 hours.

The compound of formula 7 can be prepared by using a reaction known to a skilled person. For example, the compound of formula 7 can be prepared from a

compound of formula 3 according to the following Scheme 3c in a reaction condition known to a skilled person.

**Scheme 3c:**



5

In the above Schemes from 1a to 3c, examples of suitable solvents include a mixture of any two or more of those solvents described in each Step.

The compounds of formula (I), and the intermediates above-mentioned preparation methods can be isolated and purified by conventional procedures, such as distillation, recrystallization or chromatographic purification.

The optically active compounds of this invention can be prepared by several methods. For example, the optically active compounds of this invention may be obtained by chromatographic separation, enzymatic resolution or fractional crystallization from the final compounds.

Several compounds of this invention possess an asymmetric center. Hence, the compounds can exist in separated (+)- and (-)-optically active forms, as well as in racemic one thereof. The present invention includes all such forms within its scope. Individual isomers can be obtained by known methods, such as optically selective reaction or chromatographic separation in the preparation of the final product or its intermediate.

The subject invention also includes isotopically-labelled compounds, which are identical to those recited in formula (I), but for the fact that one or more atoms are replaced by an atom having an atomic mass or mass number different from the atomic

mass or mass number usually found in nature. Examples of isotopes that can be incorporated into compounds of the invention include isotopes of hydrogen, carbon, nitrogen, oxygen, phosphorus, fluorine and chlorine, such as  $^2\text{H}$ ,  $^3\text{H}$ ,  $^{13}\text{C}$ ,  $^{14}\text{C}$ ,  $^{15}\text{N}$ ,  $^{18}\text{O}$ ,  $^{17}\text{O}$ ,  $^{31}\text{P}$ ,  $^{32}\text{P}$ ,  $^{35}\text{S}$ ,  $^{18}\text{F}$ , and  $^{36}\text{Cl}$ , respectively. Compounds of the present invention, prodrugs thereof, pharmaceutically acceptable esters of said compounds and pharmaceutically acceptable salts of said compounds, of said esters or of said prodrugs which contain the aforementioned isotopes and/or other isotopes of other atoms are within the scope of this invention. Certain isotopically-labelled compounds of the present invention, for example those into which radioactive isotopes such as  $^3\text{H}$  and  $^{14}\text{C}$  are incorporated, are useful in drug and/or substrate tissue distribution assay. Tritiated, i.e.,  $^3\text{H}$ , and carbon-14, i.e.,  $^{14}\text{C}$ , isotopes are particularly preferred for their ease of presentation and detectability. Further, substitution with heavier isotopes such as deuterium, i.e.,  $^2\text{H}$ , can afford therapeutic advantage resulting from greater metabolic stability, for example increased *in vivo* half-life or reduced dosage requirement and, hence, may be preferred in some circumstances. Isotopically labeled compounds of formula (I) of this invention and prodrugs thereof can generally be prepared by carrying out the procedure disclosed in above-disclosed Schemes and/or Examples and Preparations below, by submitting a readily available isotopically labeled reagent for a non-isotopically labeled reagent.

The present invention includes salt forms of the compounds (I) as obtained.

Pharmaceutically acceptable salts of the compounds of formula (I) include the acid addition and base addition salts (including disalts) thereof.

Pharmaceutically acceptable non-toxic salts of compounds of formula (I) may be prepared by conventional techniques by, for example, contacting said compound with a stoichiometric amount of an appropriate alkali or alkaline earth metal (sodium, potassium, calcium and magnesium) hydroxide or alkoxide in water or an appropriate organic solvent such as ethanol, isopropanol, mixtures thereof, or the like.

The bases which are used to prepare the pharmaceutically acceptable base addition salts of the acidic compounds of this invention of formula (I) are those which form non-toxic base addition salts, i.e., salts containing pharmaceutically acceptable cations, such as adenine, arginine, cytosine, lysine, benethamine (i.e., N-benzyl-2-

phenylethylamine), benzathine (i.e., N,N-dibenzylethylenediamine), choline, diolamine (i.e., diethanolamine), ethylenediamine, glucosamine, glycine, guanidine, guanine, meglumine (i.e., N-methylglucamine), nicotinamide, olamine (i.e., ethanolamine), ornithine, procaine, proline, pyridoxine, serine, tyrosine, valine and tromethamine (i.e., 5 tris or tris(hydroxymethyl)aminomethane). The base addition salts can be prepared by conventional procedures.

Insofar as the certain compounds of this invention are basic compounds, they are capable of forming a wide variety of different salts with various inorganic and organic acids.

10 The acids which are used to prepare the pharmaceutically acceptable acid addition salts of the basic compounds of this invention of formula (I) are those which form non-toxic acid addition salts, i.e., salts containing pharmaceutically acceptable anions, such as the chloride, bromide, iodide, nitrate, sulfate or bisulfate, phosphate or acid phosphate, acetate, lactate, citrate or acid citrate, tartrate or bi-tartrate, succinate, 15 malate, fumarate, gluconate, saccharate, benzoate, methanesulfonate, ethanesulfonate, benzenesulfonate, p-toluenesulfonate, adipate, aspartate, camsylate, edisylate (i.e., 1,2-ethanedisulfonate), estolate (i.e., laurylsulfate), gluceptate (i.e., glucosheptonate), gluconate, 3-hydroxy-2-naphthoate, xionofoate (i.e., 1-hydroxy-2-naphthoate), isethionate, (i.e., 2-hydroxyethanesulfonate), mucate (i.e., galactarate), 2- 20 naphsylate (i.e., naphthalenesulphonate), stearate, cholate, glucuronate, glutamate, hippurate, lactobionate, lysinate, maleate, mandelate, napadisylate, nicatinate, polygalacturonate, salicylate, sulphosalicylate, tannate, tryptophanate, borate, carbonate, oleate, phthalate and pamoate (i.e., 1,1'-methylene-bis-(2-hydroxy-3-naphthoate). Of these, we prefer edisylate (including hemi-edisylate) and 25 hydrochloride. The acid addition salts can be prepared by conventional procedures.

For a review of on suitable salts see Berge *et al.*, J. Pharm. Sci., 66, 1-19, 1977.

The present invention includes salt forms of the compounds of formula (2-A') as obtained.

Compounds of formula (2-A') may be capable of forming cations. Cations of 30 compounds of formula (2-A') may be prepared by conventional techniques by, for example, contacting said compound with a stoichiometric amount of an appropriate alkali or alkaline earth metal (sodium, potassium, calcium and magnesium) hydroxide

or alkoxide in water or an appropriate organic solvent such as ethanol, isopropanol, mixtures thereof, or the like.

The bases used to prepare the base addition salts of the acidic compounds of formula (2-A') are those which form base addition salts. Such base addition salts  
5 include pharmaceutically acceptable base addition salts as described above and salts containing cations, such as triethylamine, pyridine and ammonia.

The compounds of formula (2-A') are capable of forming a wide variety of different salts with various inorganic and organic acids.

The acids used to prepare the acid addition salts of the compound of formula  
10 (2-A') are those which form acid addition salts. Such acid addition salts include pharmaceutically acceptable acid addition salts as described above and salts containing anions, such as cyanide.

Also included within the scope of this invention are bioprecursors (also called pro-drugs) of the compounds of the formula (I). A bioprecursor of a compound of  
15 the formula (I) is a chemical derivative thereof which is readily converted back into the parent compound of the formula (I) in biological systems. In particular, a bioprecursor of a compound of the formula (I) is converted back to the parent compound of the formula (I) after the bioprecursor has been administered to, and absorbed by, a mammalian subject, e.g., a human subject. For example, it is possible  
20 to make a bioprecursor of the compounds of formula (I) in which one or both of L and W include hydroxy groups by making an ester of the hydroxy group. When only one of L and W includes hydroxy group, only mono-ester is possible. When both L and W include hydroxy, mono- and di-esters (which can be the same or different) can be made. Typical esters are simple alkanoate esters, such as acetate, propionate, butyrate,  
25 etc. In addition, when L or W includes a hydroxy group, bioprecursors can be made by converting the hydroxy group to an acyloxymethyl derivative (e.g., a pivaloyloxymethyl derivative) by reaction with an acyloxymethyl halide (e.g., pivaloyloxymethyl chloride).

When the compounds of the formula (I) of this invention may form solvates  
30 such as hydrates, such solvates are included within the scope of this invention.

Compounds of formula (I) containing one or more asymmetric carbon atoms can exist as two or more stereoisomers. Where a compound of formula (I) contains an



alkenyl or alkenylene group, geometric cis/trans (or Z/E) isomers are possible. Where the compound contains, for example, a keto or oxime group or an aromatic moiety, tautomeric isomerism ("tautomerism") can occur. It follows that a single compound may exhibit more than one type of isomerism.

- 5 Included within the scope of the present invention are all stereoisomers, geometric isomers and tautomeric forms of the compounds of formula (I), including compounds exhibiting more than one type of isomerism, and mixtures of one or more thereof. Also included are acid addition or base salts wherein the counterion is optically active, for example, D-lactate or L-lysine, or racemic, for example, DL-tartrate or DL-arginine.
- 10

Cis/trans isomers may be separated by conventional techniques well known to those skilled in the art, for example, chromatography and fractional crystallisation.

- Conventional techniques for the preparation/isolation of individual enantiomers include chiral synthesis from a suitable optically pure precursor or resolution of the racemate (or the racemate of a salt or derivative) using, for example, chiral high pressure liquid chromatography (HPLC).
- 15

- Alternatively, the racemate (or a racemic precursor) may be reacted with a suitable optically active compound, for example, an alcohol, or, in the case where the compound of formula (I) contains an acidic or basic moiety, an acid or base such as tartaric acid or 1-phenylethylamine. The resulting diastereomeric mixture may be separated by chromatography and/or fractional crystallization and one or both of the diastereoisomers converted to the corresponding pure enantiomer(s) by means well known to a skilled person.
- 20

- Stereoisomeric conglomerates may be separated by conventional techniques known to those skilled in the art - see, for example, "Stereochemistry of Organic Compounds" by E L Eliel (Wiley, New York, 1994).
- 25

- Compounds of the invention intended for pharmaceutical use may be administered as crystalline or amorphous products. They may be obtained, for example, as solid plugs, powders, or films by methods such as precipitation, crystallization, freeze drying, spray drying, or evaporative drying. Microwave or radio frequency drying may be used for this purpose.
- 30

They may be administered alone or in combination with one or more other

compounds of the invention or in combination with one or more other drugs (or as any combination thereof). Generally, they will be administered as a formulation in association with one or more pharmaceutically acceptable excipients. The term “excipient” is used herein to describe any ingredient other than the compound(s) of the invention. The choice of excipient will to a large extent depend on factors such as the particular mode of administration, the effect of the excipient on solubility and stability, and the nature of the dosage form.

Pharmaceutical compositions suitable for the delivery of compounds of the present invention and methods for their preparation will be readily apparent to those skilled in the art. Such compositions and methods for their preparation may be found, for example, in ‘Remington’s Pharmaceutical Sciences’, 19th Edition (Mack Publishing Company, 1995).

#### ORAL ADMINISTRATION

The compounds of the invention may be administered orally. Oral administration may involve swallowing, so that the compound enters the gastrointestinal tract, or buccal or sublingual administration may be employed by which the compound enters the blood stream directly from the mouth.

Formulations suitable for oral administration include solid formulations such as tablets, capsules containing particulates, liquids, or powders, lozenges (including liquid-filled), chews, multi- and nano-particulates, gels, solid solution, liposome, films (including muco-adhesive), ovules, sprays and liquid formulations.

Liquid formulations include suspensions, solutions, syrups and elixirs. Such formulations may be employed as fillers in soft or hard capsules and typically comprise a carrier, for example, water, ethanol, polyethylene glycol, propylene glycol, methylcellulose, or a suitable oil, and one or more emulsifying agents and/or suspending agents. Liquid formulations may also be prepared by the reconstitution of a solid, for example, from a sachet.

The compounds of the invention may also be used in fast-dissolving, fast-disintegrating dosage forms such as those described in Expert Opinion in Therapeutic Patents, 11 (6), 981-986 by Liang and Chen (2001).

For tablet dosage forms, depending on dose, the drug may make up from 1 wt%

to 80 wt% of the dosage form, more typically from 5 wt% to 60 wt% of the dosage form. In addition to the drug, tablets generally contain a disintegrant. Examples of disintegrants include sodium starch glycolate, sodium carboxymethyl cellulose, calcium carboxymethyl cellulose, croscarmellose sodium, crospovidone, 5 polyvinylpyrrolidone, methyl cellulose, microcrystalline cellulose, lower alkyl-substituted hydroxypropyl cellulose, starch, pregelatinised starch and sodium alginate. Generally, the disintegrant will comprise from 1 wt% to 25 wt%, preferably from 5 wt% to 20 wt% of the dosage form.

Binders are generally used to impart cohesive qualities to a tablet formulation.

10 Suitable binders include microcrystalline cellulose, gelatin, sugars, polyethylene glycol, natural and synthetic gums, polyvinylpyrrolidone, pregelatinised starch, hydroxypropyl cellulose and hydroxypropyl methylcellulose. Tablets may also contain diluents, such as lactose (monohydrate, spray-dried monohydrate, anhydrous and the like), mannitol, xylitol, dextrose, sucrose, sorbitol, microcrystalline cellulose, starch 15 and dibasic calcium phosphate dihydrate.

Tablets may also optionally comprise surface active agents, such as sodium lauryl sulfate and polysorbate 80, and glidants such as silicon dioxide and talc. When present, surface active agents may comprise from 0.2 wt% to 5 wt% of the tablet, and glidants may comprise from 0.2 wt% to 1 wt% of the tablet.

20 Tablets also generally contain lubricants such as magnesium stearate, calcium stearate, zinc stearate, sodium stearyl fumarate, and mixtures of magnesium stearate with sodium lauryl sulphate. Lubricants generally comprise from 0.25 wt% to 10 wt%, preferably from 0.5 wt% to 3 wt% of the tablet.

Other possible ingredients include anti-oxidants, colourants, flavouring agents, 25 preservatives and taste-masking agents.

Exemplary tablets contain up to about 80% drug, from about 10 wt% to about 90 wt% binder, from about 0 wt% to about 85 wt% diluent, from about 2 wt% to about 10 wt% disintegrant, and from about 0.25 wt% to about 10 wt% lubricant.

Tablet blends may be compressed directly or by roller to form tablets. Tablet 30 blends or portions of blends may alternatively be wet-, dry-, or melt-granulated, melt congealed, or extruded before tableting. The final formulation may comprise one or more layers and may be coated or uncoated; it may even be encapsulated.

The formulation of tablets is discussed in "Pharmaceutical Dosage Forms: Tablets, Vol. 1", by H. Lieberman and L. Lachman, Marcel Dekker, N.Y., N.Y., 1980 (ISBN 0-8247-6918-X).

5 Solid formulations for oral administration may be formulated to be immediate and/or modified release. Modified release formulations include delayed-, sustained-, pulsed-, controlled-, targeted and programmed release.

Suitable modified release formulations for the purposes of the invention are described in US Patent No. 6,106,864. Details of other suitable release technologies such as high energy dispersions and osmotic and coated particles are to be found in  
10 Verma *et al*, Pharmaceutical Technology On-line, 25(2), 1-14 (2001). The use of chewing gum to achieve controlled release is described in WO 00/35298.

#### PARENTERAL ADMINISTRATION

The compounds of the invention may also be administered directly into the  
15 blood stream, into muscle, or into an internal organ. Suitable means for parenteral administration include intravenous, intraarterial, intraperitoneal, intrathecal, intraventricular, intraurethral, intrasternal, intracranial, intramuscular and subcutaneous. Suitable devices for parenteral administration include needle (including microneedle) injectors, needle-free injectors and infusion techniques.

20 Parenteral formulations are typically aqueous solutions which may contain excipients such as salts, carbohydrates and buffering agents (preferably to a pH of from 3 to 9), but, for some applications, they may be more suitably formulated as a sterile non-aqueous solution or as a dried form to be used in conjunction with a suitable vehicle such as sterile, pyrogen-free water.

25 The preparation of parenteral formulations under sterile conditions, for example, by lyophilisation, may readily be accomplished using standard pharmaceutical techniques well known to those skilled in the art.

The solubility of compounds of formula (I) used in the preparation of parenteral solutions may be increased by the use of appropriate formulation techniques, such as  
30 the incorporation of solubility-enhancing agents.

Formulations for parenteral administration may be formulated to be immediate and/or modified release. Modified release formulations include delayed-, sustained-,

pulsed-, controlled-, targeted and programmed release. Thus compounds of the invention may be formulated as a solid, semi-solid, or thixotropic liquid for administration as an implanted depot providing modified release of the active compound. Examples of such formulations include drug-coated stents and PGLA  
5 microspheres.

#### TOPICAL ADMINISTRATION

The compounds of the invention may also be administered topically to the skin or mucosa, that is, dermally or transdermally. Typical formulations for this purpose  
10 include gels, hydrogels, lotions, solutions, creams, ointments, dusting powders, dressings, foams, films, skin patches, wafers, implants, sponges, fibres, bandages and microemulsions. Liposomes may also be used. Typical carriers include alcohol, water, mineral oil, liquid petrolatum, white petrolatum, glycerin, polyethylene glycol and propylene glycol. Penetration enhancers may be incorporated - see, for example, J  
15 Pharm Sci, 88 (10), 955-958 by Finnin and Morgan (October 1999).

Other means of topical administration include delivery by electroporation, iontophoresis, phonophoresis, sonophoresis and microneedle or needle-free (e.g. Powderject™, Bioject™, etc.) injection.

Formulations for topical administration may be formulated to be immediate  
20 and/or modified release. Modified release formulations include delayed-, sustained-, pulsed-, controlled-, targeted and programmed release.

#### INHALED/INTRANASAL ADMINISTRATION

The compounds of the invention can also be administered intranasally or by  
25 inhalation, typically in the form of a dry powder (either alone, as a mixture, for example, in a dry blend with lactose, or as a mixed component particle, for example, mixed with phospholipids, such as phosphatidylcholine) from a dry powder inhaler or as an aerosol spray from a pressurised container, pump, spray, atomiser (preferably an atomiser using electrohydrodynamics to produce a fine mist), or nebuliser, with or  
30 without the use of a suitable propellant, such as 1,1,1,2-tetrafluoroethane or 1,1,1,2,3,3,3-heptafluoropropane. For intranasal use, the powder may comprise a bioadhesive agent, for example, chitosan or cyclodextrin.

The pressurised container, pump, spray, atomizer, or nebuliser contains a solution or suspension of the compound(s) of the invention comprising, for example, ethanol, aqueous ethanol, or a suitable alternative agent for dispersing, solubilising, or extending release of the active, a propellant(s) as solvent and an optional surfactant, such as sorbitan trioleate, oleic acid, or an oligolactic acid.

Prior to use in a dry powder or suspension formulation, the drug product is micronised to a size suitable for delivery by inhalation (typically less than 5 microns). This may be achieved by any appropriate comminuting method, such as spiral jet milling, fluid bed jet milling, supercritical fluid processing to form nanoparticles, high pressure homogenisation, or spray drying.

Capsules (made, for example, from gelatin or HPMC), blisters and cartridges for use in an inhaler or insufflator may be formulated to contain a powder mix of the compound of the invention, a suitable powder base such as lactose or starch and a performance modifier such as *L*-leucine, mannitol, or magnesium stearate. The lactose may be anhydrous or in the form of the monohydrate, preferably the latter. Other suitable excipients include dextran, glucose, maltose, sorbitol, xylitol, fructose, sucrose and trehalose.

A suitable solution formulation for use in an atomiser using electrohydrodynamics to produce a fine mist may contain from 1 $\mu$ g to 20mg of the compound of the invention per actuation and the actuation volume may vary from 1 $\mu$ l to 100 $\mu$ l. A typical formulation may comprise a compound of formula (I), propylene glycol, sterile water, ethanol and sodium chloride. Alternative solvents which may be used instead of propylene glycol include glycerol and polyethylene glycol.

Suitable flavours, such as menthol and levomenthol, or sweeteners, such as saccharin or saccharin sodium, may be added to those formulations of the invention intended for inhaled/intranasal administration.

Formulations for inhaled/intranasal administration may be formulated to be immediate and/or modified release using, for example, poly(DL-lactic-coglycolic acid (PLGA). Modified release formulations include delayed-, sustained-, pulsed-, controlled-, targeted and programmed release.

In the case of dry powder inhalers and aerosols, the dosage unit is determined by means of a valve which delivers a metered amount. Units in accordance with the

invention are typically arranged to administer a metered dose or "puff" containing from 1 to 100  $\mu\text{g}$  of the compound of formula (I). The overall daily dose will typically be in the range 50  $\mu\text{g}$  to 20 mg which may be administered in a single dose or, more usually, as divided doses throughout the day.

5

#### RECTAL/INTRAVAGINAL ADMINISTRATION

The compounds of the invention may be administered rectally or vaginally, for example, in the form of a suppository, pessary, or enema. Cocoa butter is a traditional suppository base, but various alternatives may be used as appropriate.

10

Formulations for rectal/vaginal administration may be formulated to be immediate and/or modified release. Modified release formulations include delayed-, sustained-, pulsed-, controlled-, targeted and programmed release.

#### 15 OCULAR/AURAL ADMINISTRATION

The compounds of the invention may also be administered directly to the eye or ear, typically in the form of drops of a micronised suspension or solution in isotonic, pH-adjusted, sterile saline. Other formulations suitable for ocular and aural administration include ointments, biodegradable (e.g. absorbable gel sponges, collagen) and non-biodegradable (e.g. silicone) implants, wafers, lenses and particulate or vesicular systems, such as niosomes or liposomes. A polymer such as crossed-linked polyacrylic acid, polyvinylalcohol, hyaluronic acid, a cellulosic polymer, for example, hydroxypropylmethylcellulose, hydroxyethylcellulose, or methyl cellulose, or a heteropolysaccharide polymer, for example, gelatin gum, may be incorporated together with a preservative, such as benzalkonium chloride. Such formulations may also be delivered by iontophoresis.

Formulations for ocular/aural administration may be formulated to be immediate and/or modified release. Modified release formulations include delayed-, sustained-, pulsed-, controlled-, targeted, or programmed release.

30

#### OTHER TECHNOLOGIES

The compounds of the invention may be combined with soluble

macromolecular entities, such as cyclodextrin and suitable derivatives thereof or polyethylene glycol-containing polymers, in order to improve their solubility, dissolution rate, taste-masking, bioavailability and/or stability for use in any of the aforementioned modes of administration.

5

Drug-cyclodextrin complexes, for example, are found to be generally useful for most dosage forms and administration routes. Both inclusion and non-inclusion complexes may be used. As an alternative to direct complexation with the drug, the cyclodextrin may be used as an auxiliary additive, *i.e.* as a carrier, diluent, or solubiliser. Most commonly used for these purposes are alpha-, beta- and gamma-cyclodextrins, examples of which may be found in International Patent Applications Nos. WO 91/11172, WO 94/02518 and WO 98/55148.

10

#### KIT-OF-PARTS

Inasmuch as it may desirable to administer a combination of active compounds, for example, for the purpose of treating a particular disease or condition, it is within the scope of the present invention that two or more pharmaceutical compositions, at least one of which contains a compound in accordance with the invention, may conveniently be combined in the form of a kit suitable for coadministration of the compositions.

20

Thus the kit of the invention comprises two or more separate pharmaceutical compositions, at least one of which contains a compound of formula (I) in accordance with the invention, and means for separately retaining said compositions, such as a container, divided bottle, or divided foil packet. An example of such a kit is the familiar blister pack used for the packaging of tablets, capsules and the like.

25

The kit of the invention is particularly suitable for administering different dosage forms, for example, oral and parenteral, for administering the separate compositions at different dosage intervals, or for titrating the separate compositions against one another. To assist compliance, the kit typically comprises directions for administration and may be provided with a so-called memory aid.

30

#### DOSAGE



For administration to human patients, the total daily dose of the compounds of the invention is typically in the range 0.05 mg to 100 mg depending, of course, on the mode of administration, preferred in the range 0.1 mg to 50 mg and more preferred in the range 0.5 mg to 20 mg. For example, oral administration may require a total daily  
5 dose of from 1 mg to 20 mg, while an intravenous dose may only require from 0.5 mg to 10 mg. The total daily dose may be administered in single or divided doses.

These dosages are based on an average human subject having a weight of about 65kg to 70kg. The physician will readily be able to determine doses for subjects whose weight falls outside this range, such as infants and the elderly.

10 For the avoidance of doubt, references herein to "treatment" include references to curative, palliative and prophylactic treatment.

A 5-HT<sub>4</sub> agonist of the present invention may be usefully combined with another pharmacologically active compound, or with two or more other  
15 pharmacologically active compounds, particularly in the treatment of gastroesophageal reflux disease. For example, a 5-HT<sub>4</sub> agonist, particularly a compound of the formula (I), or a pharmaceutically acceptable salt or solvate thereof, as defined above, may be administered simultaneously, sequentially or separately in combination with one or more agents selected from:

- 20 (i) histamine H<sub>2</sub> receptor antagonists, e.g. ranitidine, lafutidine, nizatidine, cimetidine, famotidine and roxatidine;
- (ii) proton pump inhibitors, e.g. omeprazole, esomeprazole, pantoprazole, rabeprazole, tenatoprazole, ilaprazole and lansoprazole;
- (iii) Acid pump antagonists, e.g. soraprazan, revaprazan(YH-1885), AZD-0865,  
25 CS-526, AU-2064 and YJA-20379-8;
- (iv) oral antacid mixtures, e.g. Maalox<sup>®</sup>, Aludrox<sup>®</sup> and Gaviscon<sup>®</sup>;
- (v) mucosal protective agents, e.g. polaprezinc, ecabet sodium, rebamipide, teprenone, cetraxate, sucralfate, chloropylline-copper and plaunotol;
- (vi) GABA<sub>B</sub> agonists, e.g. baclofen and AZD-3355;
- 30 (vii)  $\alpha$ 2 agonists, e.g. clonidine, medetomidine, lofexidine, moxonidine, tizanidine, guanfacine, guanabnz, talipexole and dexmedetomidine;
- (viii) Xanthin derivatives, e.g. Theophylline, aminophylline and doxofylline;

- (ix) calcium channel blockers, e.g. amlodipine, lacidipine, flutopidine, amlodipine, lomerizine, diltiazem, gallopamil, efonidipine, nisoldipine, amlodipine, lercanidipine, bevantolol, nicardipine, isradipine, benidipine, verapamil, nitrendipine, barnidipine, propafenone, manidipine, bepridil, nifedipine, nilvadipine, nimodipine, nifedipine and fasudil;
- 5 (x) benzodiazepine agonists, e.g. diazepam, zaleplon, zolpidem, haloxazolam, clonazepam, prazepam, quazepam, flutazolam, triazolam, lormetazepam, midazolam, tofisopam, clobazam, flunitrazepam and flutoprazepam;
- (xi) prostaglandin analogues, e.g. Prostaglandin, misoprostol, treprostinil, 10 esoprostenoil, latanoprost, iloprost, beraprost, enprostil, ibudilast and ozagrel;
- (xii) histamine H<sub>3</sub> agonists, e.g. R- $\alpha$ -methylhistamine and BP-294;
- (xiii) anti-gastric agents, e.g. Anti-gastrin vaccine, itriglumide and Z-360;
- (xiv) 5-HT<sub>3</sub> antagonists, e.g. dolasetron, palonosetron, alosetron, azasetron, ramosetron, mitrazapine, granisetron, tropisetron, E-3620, ondansetron and 15 indisetron;
- (xv) tricyclic antidepressants, e.g. imipramine, amitriptyline, clomipramine, amoxapine and lofepramine;
- (xvi) GABA agonists, e.g. gabapentin, topiramate, cinolazepam, clonazepam, progabide, brotizolam, zopiclone, pregabalin and eszopiclone;
- 20 (xvii) opioid analgesics, e.g. morphine, heroin, hydromorphone, oxymorphone, levorphanol, levallorphan, methadone, meperidine, fentanyl, cocaine, codeine, dihydrocodeine, oxycodone, hydrocodone, propoxyphene, nalmefene, nalorphine, naloxone, naltrexone, buprenorphine, butorphanol, nalbuphine and pentazocine;
- (xviii) somatostatin analogues, e.g. octreotide, AN-238 and PTR-3173;
- 25 (xix) Cl Channel activator: e.g. lubiprostone;
- (xx) selective serotonin reuptake inhibitors, e.g. sertraline, escitalopram, fluoxetine, nefazodone, fluvoxamine, citalopram, milnacipran, paroxetine, venlafaxine, tramadol, sibutramine, duloxetine, desvenlafaxine and depocoxetine;
- (xxi) anticholinergics, e.g. dicyclomine and hyoscyamine;
- 30 (xxii) laxatives, e.g. Trifyba<sup>®</sup>, Fybogel<sup>®</sup>, Konsyl<sup>®</sup>, Isogel<sup>®</sup>, Regular<sup>®</sup>, Celevac<sup>®</sup> and Normacol<sup>®</sup>;
- (xxiii) fiber products, e.g. Metamucil<sup>®</sup>;

- (xxiv) antispasmodics, e.g.: mebeverine;
- (xxv) dopamine antagonists, e.g. metoclopramide, domperidone and levosulpiride;
- (xxvi) cholinergics, e.g. neostigmine
- (xxvii) AChE inhibitor: galantamine, metrifonate, rivastigmine, itopride and donepezil;
- (xxviii) Tachykinin (NK) antagonists, particularly NK-3, NK-2 and NK-1 e.g. antagonists, nepadutant, saredutant, talnetant, ( $\alpha$ R,9R)-7-[3,5-bis(trifluoromethyl)benzyl]-8,9,10,11-tetrahydro-9-methyl-5-(4-methylphenyl)-7H-[1,4]diazocino[2,1-g][1,7]naphthridine-6-13-dione (TAK-637), 5-[[[(2R,3S)-2-[(1R)-1-[3,5-bis(trifluoromethyl)phenyl]ethoxy-3-(4-fluorophenyl)-4-morpholinyl]methyl]-1,2-dihydro-3H-1,2,4-triazol-3-one (MK-869), lanepitant, dapitant and 3-[[[2-methoxy-5-(trifluoromethoxy)phenyl]methylamino]-2-phenyl-piperidine (2S,3S).

#### Method for assessing biological activities:

- 15 The 5-HT<sub>4</sub> receptor binding affinities of the compounds of this invention are determined by the following procedures.

#### Membrane Preparation

- Pig heads were supplied from an abattoir. Striatal tissues were dissected, weighed and homogenized in 15 volumes of 50 mM ice-cold HEPES (pH 7.5) in a Polytron homogenizer (30 sec at full speed). Suspension was centrifuged at 48,000g and 4°C for 15 min. The resulting pellet was resuspended in an appropriate volume of 50 mM ice-cold HEPES, dispensed into aliquots and stored at -80°C until use.

- Bovine heads were also supplied from an abattoir. Striatal tissues were dissected, weighed and homogenized in 20 volumes of 50 mM ice-cold Tris-HCl (pH 7.4) in a Polytron homogenizer (30 sec at full speed). Suspension was centrifuged at 20,000g and 4°C for 30 min. The resulting pellet was resuspended in 15 volumes of 50 mM ice-cold Tris-HCl, homogenized and centrifuged again in the same way. The final pellet was resuspended in an appropriate volume of 50 mM Tris-HCl, dispensed into aliquots and stored at -80°C until use.

- 30 Cerebral cortical tissues were removed from male Sprague-Dawley (SD) rats (Japan SLC), weighed and placed in 10 volumes of 50 mM ice-cold Tris-HCl (pH 7.5). This was homogenized in a Polytron homogenizer (30 sec at full speed) and

subsequently centrifuged at 48,000g and 4°C for 15 min. The resulting pellet was resuspended in 50 mM ice-cold Tris-HCl, homogenized and centrifuged again in the same way. The final pellet was resuspended in an appropriate volume of 50 mM Tris-HCl, dispensed into aliquots and stored at -80°C until use.

- 5       The protein concentrations of homogenates were determined by Bradford method or BCA protein method (Pierce) with BSA as a standard.

### **Binding assays**

- Affinity of compounds for pig or bovine 5-HT<sub>4</sub> and rat 5-HT<sub>3</sub> receptors were assessed with using radiolabeled specific ligands, GR 113808 ({1-[2-(methylsulfonyl)ethyl]-4-piperidinyl}[methyl-<sup>3</sup>H]-1*H*-indole-3-carboxylate) and BRL 43694 (1-Methyl-*N*-(9-[methyl-<sup>3</sup>H]-9-azabicyclo[3.3.1]non-3-yl)-1*H*-indazole-3-carboxamide). Compounds were incubated with 25-100 pM of [<sup>3</sup>H]-GR 113808 (Amersham) and 0.6-1 mg protein of pig or bovine striatal membranes suspended in a final volume of 0.8-1 ml of 50 mM Tris-HCl (pH 7.5). Nonspecific binding was determined with 10-50 μM 5-HT. The binding of 0.3 nM [<sup>3</sup>H]-BRL 43694 (NEN) was measured using 400 μg protein of rat cortical membranes suspended in a final volume of 500 μl of 50 mM Tris-HCl (pH 7.5). Nonspecific binding was determined with 10 μM 5-HT.
- 10  
15

- The plates were incubated at room temperature on a plate shaker for 30min.
- 20   The assays were stopped by rapid filtration using a Brandell cell harvester through Wallace-B filters pre-soaked in 0.2% poly(ethylenimine) at 4°C for 60-90min. The filters were washed three times with 1 ml of ice-cold 50 mM HEPES, and were dried in a microwave or at room temperature. They were bagged and heated with meltilex scintillant (Wallace) or soaked in BetaplateScint (Wallace). Receptor-bound
- 25   radioactivity was quantified using Big-spot counter, Betaplate counter (Wallace) or LS counter (Packard).

### **Human 5-HT<sub>4</sub> binding(1)**

- Human 5-HT<sub>4(d)</sub> transfected HEK293 cells were prepared and grown in-house. The collected cells were suspended in 50 mM HEPES (pH 7.4 at 4°C) supplemented with protease inhibitor cocktail (Boehringer, 1:1000 dilution) and homogenized using a hand held Polytron PT 1200 disruptor set at full power for 30 sec on ice. The homogenates were centrifuged at 40,000 x g at 4 °C for 30 min. The pellets were then
- 30

resuspended in 50 mM HEPES (pH 7.4 at 4 °C) and centrifuged once more in the same manner. The final pellets were resuspended in an appropriate volume of 50 mM HEPES (pH 7.4 at 25 °C), homogenized, aliquoted and stored at -80°C until use. An aliquot of membrane fractions was used for protein concentration determination using

- 5 BCA protein assay kit (PIERCE) and ARVOsx plate reader (Wallac).

For the binding experiments, 25 µl of test compounds were incubated with 25 µl of [<sup>3</sup>H]-GR113808 (Amersham, final 0.2 nM) and 150 µl of membrane homogenate and WGA-SPA beads (Amersham) suspension solutions (10 µg protein and 1mg SPA beads/well) for 60 minutes at room temperature. Nonspecific binding  
10 was determined by 1 µM GR113808 (Tocris) at the final concentration. Incubation was terminated by centrifugation at 1000 rpm. Receptor-bound radioactivity was quantified by counting with MicroBeta plate counter (Wallac).

All compounds prepared in the working examples as described below were tested by this method, and they showed K<sub>i</sub> values from 0.3 nM to 30 nM with respect  
15 to inhibition of binding at the 5-HT<sub>4</sub> receptor.

#### **Human 5-HT<sub>4</sub> binding(2)**

Human 5-HT<sub>4(d)</sub> transfected HEK293 cells were prepared and grown in-house. The collected cells were suspended in 50 mM Tris buffer (pH 7.4 at 4°C) supplemented with protease inhibitor cocktail (Boehringer, 1:1000 dilution) and  
20 homogenized using a hand held Polytron PT 1200 disruptor set at full power for 30 sec on ice. The homogenates were centrifuged at 40,000 x g at 4 °C for 10 min. The pellets were then resuspended in 50 mM Tris buffer (pH 7.4 at 4 °C) and centrifuged once more in the same manner. The final pellets were resuspended in an appropriate volume of 50 mM Tris buffer (pH 7.4 at 25 °C) containing 10 mM MgCl<sub>2</sub>,  
25 homogenized, aliquoted and stored at -80°C until use. An aliquot of membrane fractions was used for protein concentration determination using BCA protein assay kit (PIERCE) and ARVOsx plate reader (Wallac).

For the binding experiments, 50 µl of test compounds were incubated with 50 µl of [<sup>3</sup>H] 5-HT (Amersham, final 8.0 nM) and 400 µl of membrane homogenate (300 µg  
30 protein/ tube) for 60 minutes at room temperature. Nonspecific binding was determined by 50 µM GR113808 (Tocris) at the final concentration. All incubations were terminated by rapid vacuum filtration over 0.2 % PEI soaked glass fiber filter

papers using BRANDEL harvester followed by three washes with 50 mM Tris buffer (pH 7.4 at 25 °C). Receptor-bound radioactivity was quantified by liquid scintillation counting using Packard LS counter.

All compounds of Examples showed 5HT<sub>4</sub> receptor affinity.

5

#### Functional Assay:

The presence of 5-HT<sub>4</sub> receptors in the rat oesophagus and the ability to demonstrate partial agonism in the TMM preparation are reported in the literature (See G.S. Baxter et al. Naunyn-Schmiedeberg's Arch Pharmacol (1991) 343: 439-446; M. Yukiko et al. JPET (1997) 283: 1000-1008; and J.J. Reeves et al. Br. J. Pharmacol. (1991) 103: 1067-1072). More specifically, partial agonist activity can be measured according to the following procedures.

Male SD rats (Charles River) weighing 250-350g were stunned and then killed by cervical dislocation. The oesophagus was dissected from immediately proximal to the stomach (including piece of stomach to mark distal end) up to the level of the trachea and then placed in fresh Krebs' solution.

The outer skeletal muscle layer was removed in one go by peeling it away from the underlying smooth muscle layer using forceps (stomach to tracheal direction). The remaining inner tube of smooth muscle was known as the TMM. This was trimmed to 2 cm from the original 'stomach-end' and the rest discarded.

The TMMs were mounted as whole 'open' tubes in longitudinal orientation in 5ml organ baths filled with warm (32°C) aerated Krebs. Tissues were placed under an initial tension of 750mg and allowed to equilibrate for 60 minutes. The tissues were re-tensioned twice at 15 minute intervals during the equilibration period. The pump flow rate was set to 2ml / min during this time.

Following equilibration, the pump was switched off. The tissues were exposed to 1  $\mu$  M carbachol and contracted and reached a steady contractile plateau within 15 minutes. Tissues were then subject to 1  $\mu$  M 5-HT (this was to prime the tissues). The tissues relaxed in response to 5-HT fairly rapidly - within 1 minute. As soon as maximal relaxation has occurred and a measurement taken, the tissues were washed at maximum rate (66ml/min) for at least 1 minute and until the original baseline (pre-carbachol and 5-HT) has returned (usually, the baseline drops below the

original one following initial equilibration). The pump flow rate was reduced to 2ml/min and the tissues left for 60 minutes.

A cumulative concentration-effect-curve (CEC) to 5-HT was constructed across the range 0.1 nM to 1  $\mu$ M, in half-log unit increments (*5-HT curve 1* for data analysis). Contact time between doses was 3 minutes or until plateau established. Tissues responded quicker as concentration of 5-HT in the bath increases. At the end of the curve, the tissues were washed (at maximum rate) as soon as possible to avoid desensitisation of receptors. Pump rate was reduced to 2ml/min and the tissues left for 60 minutes.

A second CEC was carried out - either to 5-HT (for time control tissues), another 5-HT<sub>4</sub> agonist (standard) or a test compound (*curve 2* for data analysis). Contact time varied for other 5-HT<sub>4</sub> agonists and test compounds and was tailored according to the tissues' individual responses to each particular agent. In tissues exposed to a test compound, a high concentration (1  $\mu$ M) of a 5-HT<sub>4</sub> antagonist (SB 203,186: 1*H*-Indole-3-carboxylic acid, 2-(1-piperidinyl)ethyl ester, Tocris) was added to the bath following the last concentration of test compound. This was to see if any agonist-induced relaxation (if present) could be reversed. SB 203,186 reversed 5-HT induced relaxation, restoring the tissue's original degree of carbachol-induced tone.

Agonist activity of test compounds was confirmed by pre-incubating tissues with 100nM standard 5HT<sub>4</sub> antagonist such as SB 203,186. SB 203,186 was added to the bath 5 minutes before the addition of carbachol prior to *curve 2*. Tissues must be 'paired' for data analysis i.e. the test compound in the absence of SB 203,186 in one tissue was compared with the test compound in the presence of SB 203,186 in a separate tissue. It was not possible to carry out a *curve 3* i.e. 5-HT *curve 1*, followed by the test compound *curve 2* (- SB 203,186), followed by the test compound *curve 3* (+ SB 203,186).

#### Agonist-induced cAMP elevation in human 5-HT<sub>4(4)</sub> transfected HEK293 cells

Human 5-HT<sub>4(4)</sub> transfected HEK293 cells were established in-house. The cells were grown at 37°C and 5% CO<sub>2</sub> in DMEM supplemented with 10% FCS, 20 mM HEPES (pH 7.4), 200  $\mu$ g/ml hygromycin B (Gibco), 100 units/ml penicillin and

100 µg/ml streptomycin.

The cells were grown to 60-80% confluence. On the previous day before treatment with compounds dialyzed FCS (Gibco) was substituted for normal and the cells were incubated overnight.

- 5 Compounds were prepared in 96-well plates (12.5 µl/well). The cells were harvested with PBS/1 mM EDTA, centrifuged and washed with PBS. At the beginning of the assay, cell pellet was resuspended in DMEM supplemented with 20 mM HEPES, 10 µM pargyline (Sigma) and 1 mM 3-isobutyl-1-methylxanthine (Sigma) at the concentration of  $1.6 \times 10^5$  cells/ml and left for 15 minutes at room temperature. The
- 10 reaction was initiated by addition of the cells into plates (12.5 µl/well). After incubation for 15 minutes at room temperature, 1% Triton X-100 was added to stop the reaction (25 µl/well) and the plates were left for 30 minutes at room temperature. Homogenous time-resolved fluorescence-based cAMP (Schering) detection was made according to the manufacturer's instruction. ARVOsx multilabel counter (Wallac)
- 15 was used to measure HTRF (excitation 320 nm, emission 665 nm/620 nm, delay time 50 µs, window time 400 µs).
- Data was analyzed based on the ratio of fluorescence intensity of each well at 620 nm and 665 nm followed by cAMP quantification using cAMP standard curve.
- Enhancement of cAMP production elicited by each compound was normalized to the
- 20 amount of cAMP produced by 1000 nM serotonin (Sigma).

All compounds of Examples showed 5HT<sub>4</sub> receptor agonistic activity.

### **Human dofetilide binding**

Human HERG transfected HEK293S cells were prepared and grown in-house.

- 25 The collected cells were suspended in 50 mM Tris-HCl (pH 7.4 at 4°C) and homogenized using a hand held Polytron PT 1200 disruptor set at full power for 20 sec on ice. The homogenates were centrifuged at 48,000 x g at 4 °C for 20 min. The pellets were then resuspended, homogenized, and centrifuged once more in the same manner. The final pellets were resuspended in an appropriate volume of 50 mM Tris-
- 30 HCl, 10 mM KCl, 1 mM MgCl<sub>2</sub> (pH 7.4 at 4°C), homogenized, aliquoted and stored at -80°C until use. An aliquot of membrane fractions was used for protein concentration determination using BCA protein assay kit (PIERCE) and ARVOsx



plate reader (Wallac).

Binding assays were conducted in a total volume of 200  $\mu$ l in 96-well plates. Twenty  $\mu$ l of test compounds were incubated with 20  $\mu$ l of [ $^3$ H]-dofetilide (Amersham, final 5 nM) and 160  $\mu$ l of membrane homogenate (25  $\mu$ g protein) for 60 minutes at room temperature. Nonspecific binding was determined by 10  $\mu$ M dofenilide at the final concentration. Incubation was terminated by rapid vacuum filtration over 0.5% presoaked GF/B Betaplate filter using Skatron cell harvester with 50 mM Tris-HCl, 10 mM KCl, 1 mM MgCl<sub>2</sub>, pH 7.4 at 4°C. The filters were dried, put into sample bags and filled with Betaplate Scint. Radioactivity bound to filter was counted with Wallac Betaplate counter.

#### HERG assay

HEK 293 cells which stably express the HERG potassium channel were used for electrophysiological study. The methodology for stable transfection of this channel in HEK cells can be found elsewhere (Z.Zhou et al., 1998, Biophysical journal, 74, pp230-241). Before the day of experimentation, the cells were harvested from culture flasks and plated onto glass coverslips in a standard MEM medium with 10% FCS. The plated cells were stored in an incubator at 37°C maintained in an atmosphere of 95%O<sub>2</sub>/5%CO<sub>2</sub>. Cells were studied between 15-28hrs after harvest.

HERG currents were studied using standard patch clamp techniques in the whole-cell mode. During the experiment the cells were superfused with a standard external solution of the following composition (mM); NaCl, 130; KCl, 4; CaCl<sub>2</sub>, 2; MgCl<sub>2</sub>, 1; Glucose, 10; HEPES, 5; pH 7.4 with NaOH. Whole-cell recordings was made using a patch clamp amplifier and patch pipettes which have a resistance of 1-3M $\Omega$  when filled with the standard internal solution of the following composition (mM); KCl, 130; MgATP, 5; MgCl<sub>2</sub>, 1.0; HEPES, 10; EGTA 5, pH 7.2 with KOH. Only those cells with access resistances below 15M $\Omega$  and seal resistances >1G $\Omega$  was accepted for further experimentation. Series resistance compensation was applied up to a maximum of 80%. No leak subtraction was done. However, acceptable access resistance depended on the size of the recorded currents and the level of series resistance compensation that can safely be used. Following the achievement of whole cell configuration and sufficient for cell dialysis with pipette solution (>5min),

a standard voltage protocol was applied to the cell to evoke membrane currents. The voltage protocol is as follows. The membrane was depolarized from a holding potential of -80mV to +20mV for 1000ms. This was followed by a descending voltage ramp (rate 0.5mV msec<sup>-1</sup>) back to the holding potential. The voltage protocol was applied to a cell continuously throughout the experiment every 4 seconds (0.25Hz). The amplitude of the peak current elicited around -40mV during the ramp was measured. Once stable evoked current responses were obtained in the external solution, vehicle (0.5% DMSO in the standard external solution) was applied for 10-20 min by a peristaltic pump. Provided there were minimal changes in the amplitude of the evoked current response in the vehicle control condition, the test compound of either 0.3, 1, 3, 10μM was applied for a 10 min period. The 10 min period included the time which supplying solution was passing through the tube from solution reservoir to the recording chamber via the pump. Exposing time of cells to the compound solution was more than 5min after the drug concentration in the chamber well reached the attempting concentration. There reversibility. Finally, the cells was exposed to high dose of dofetilide (5μM), a specific I<sub>Kr</sub> blocker, to evaluate the insensitive endogenous current.

All experiments were performed at room temperature (23 ± 1°C). Evoked membrane currents were recorded on-line on a computer, filtered at 500-1KHz (Bessel -3dB) and sampled at 1-2KHz using the patch clamp amplifier and a specific data analyzing software. Peak current amplitude, which occurred at around -40mV, was measured off line on the computer.

The arithmetic mean of the ten values of amplitude was calculated under control conditions and in the presence of drug. Percent decrease of I<sub>N</sub> in each experiment was obtained by the normalized current value using the following formula:  $I_N = (1 - I_D/I_C) \times 100$ , where I<sub>D</sub> is the mean current value in the presence of drug and I<sub>C</sub> is the mean current value under control conditions. Separate experiments were performed for each drug concentration or time-matched control, and arithmetic mean in each experiment is defined as the result of the study.

#### Half-life in human liver microsomes (HLM)

Test compounds (1 μM) were incubated with 3.3 mM MgCl<sub>2</sub> and 0.78

mg/mL HLM (HL101) in 100 mM potassium phosphate buffer (pH 7.4) at 37°C on the 96-deep well plate. The reaction mixture was split into two groups, a non-P450 and a P450 group. NADPH was only added to the reaction mixture of the P450 group. An aliquot of samples of P450 group was collected at 0, 10, 30, and 60 min time point, where 0 min time point indicated the time when NADPH was added into the reaction mixture of P450 group. An aliquot of samples of non-P450 group was collected at -10 and 65 min time point. Collected aliquots were extracted with acetonitrile solution containing an internal standard. The precipitated protein was spun down in centrifuge (2000 rpm, 15 min). The compound concentration in supernatant was measured by LC/MS/MS system.

The half-life value was obtained by plotting the natural logarithm of the peak area ratio of compounds/ internal standard versus time. The slope of the line of best fit through the points yields the rate of metabolism (k). This was converted to a half-life value using following equations:

$$\text{Half-life} = \ln 2 / k$$

#### Method of gastric emptying model in rats:

The effects of compounds on gastric emptying in rats were examined by the modified method of D. A. Droppleman et al. (J. Pharmacol. Methods 4, 227-230 (1980)). The test meal, non-fat caloric meal, was prepared according to the method of S. Ueki et al (Arzneim.-Forsch./Drug Res. 49 (II), 618-625 (1999)). IGS-SD rats (Male, 7w, 230-270 g) were purchased from Charles River Japan (Atsugi). These rats were used in the experiments after one week acclimatization. In the experiments, rats were fasted 15 hrs before the experiments but allowed free access to water. Forty-five minutes prior to the start of the experiment, water was removed from the cage to prevent rats from taking water. Five minutes before the test meal administration, test compounds, cisapride or vehicle were dosed via an appropriate route to rats (n=8-10) in a volume of 0.1 ml per 100 g body weight. Cisapride (3 mg/kg) was used as a positive control for the experiment. Rats were given 3 ml of the test meal by gavage and were returned to the cages. Thirty minutes after the meal administration, rats were culled by CO<sub>2</sub> exposure. Following a midline laparotomy, the stomach is ligated at the lower esophageal sphincter (LES) and pylorus. Then the stomach was removed and

weighed (A). After the stomach was opened and rinsed with 0.9 % saline, it was blotted the face with the tissue to remove any excess liquid and weighed again (B). After avoiding the rats that had eaten feces or given artificial miss, gastric emptying rate for individual animals was calculated by the formula:

- 5 GE rate (%)=(A-B)/weight of the test meal.

**Gastric motility in conscious dogs:**

The surgical operation in dogs was performed by the modified method of Z. Itoh et al. (Gastroenterol. Jpn., 12, 275-283 (1977)). The effects of test compounds on gastric  
10 motility in dogs were examined by the modified method of N. Toshida et al. (J.Pharmacol.Exp/Ther., 257, 781-787 (1991)).

An evaluation in the fasted state: Animals were chronically implanted with a strain gauge force transducer on the gastric body, and fasted overnight prior to the experiment. The gastric motility was continuously recorded by a telemetry system  
15 for 8 h after administration of the compound. To quantitate the change in gastrointestinal motility, the motor index was determined as the area under the contraction curves during each 2 h period divided by the peak height of interdigestive migrating contraction.

An evaluation in the postprandial state: Animals were chronically implanted with a strain gauge force transducer on the gastric body, and fasted overnight prior to the experiment. Postprandial motility was induced by feeding with solid meal (100 grams), and the compound was administered 2 h later. The gastric motility was continuously recorded by a telemetry system for 8 h after administration of the compound. The motor index was determined to quantitate the change in  
25 gastrointestinal motility as the area under the contraction curves during each 1 h period divided by the area under the contraction curves for 1 h before the compound administration.

The compounds of formula (I) of this invention can be administered via  
30 either the oral, parenteral or topical routes to mammals. In general, these compounds are most desirably administered to humans in doses ranging from 0.3 mg to 750 mg per day, preferably from 10 mg to 500 mg per day, although variations will

necessarily occur depending upon the weight and condition of the subject being treated, the disease state being treated and the particular route of administration chosen. However, for example, a dosage level that is in the range of from 0.06 mg to 2 mg per kg of body weight per day is most desirably employed for treatment of inflammation.

The compounds of the present invention may be administered alone or in combination with pharmaceutically acceptable carriers or diluents by either of the above routes previously indicated, and such administration can be carried out in single or multiple doses. More particularly, the novel therapeutic agents of the invention can be administered in a wide variety of different dosage forms, i.e., they may be combined with various pharmaceutically acceptable inert carriers in the form of tablets, capsules, lozenges, troches, hard candies, powders, sprays, creams, salves, suppositories, jellies, gels, pastes, lotions, ointments, aqueous suspensions, injectable solutions, elixirs, syrups, and the like. Such carriers include solid diluents or fillers, sterile aqueous media and various non-toxic organic solvents, etc. Moreover, oralpharmaceutical compositions can be suitably sweetened and/or flavored. In general, the therapeutically-effective compounds of this invention are present in such dosage forms at concentration levels ranging 5% to 70% by weight, preferably 10% to 50% by weight.

For oral administration, tablets containing various excipients such as microcrystalline cellulose, sodium citrate, calcium carbonate, dipotassium phosphate and glycine may be employed along with various disintegrants such as starch and preferably corn, potato or tapioca starch, alginic acid and certain complex silicates, together with granulation binders like polyvinylpyrrolidone, sucrose, gelatin and acacia. Additionally, lubricating agents such as magnesium stearate, sodium lauryl sulfate and talc are often very useful for tableting purposes. Solid compositions of a similar type may also be employed as fillers in gelatin capsules; preferred materials in this connection also include lactose or milk sugar as well as high molecular weight polyethylene glycols. When aqueous suspensions and/or elixirs are desired for oral administration, the active ingredient may be combined with various sweetening or flavoring agents, coloring matter or dyes, and, if so desired, emulsifying and/or suspending agents as well, together with such diluents as water, ethanol, propylene

glycol, glycerin and various like combinations thereof.

For parenteral administration, solutions of a compound of the present invention in either sesame or peanut oil or in aqueous propylene glycol may be employed. The aqueous solutions should be suitably buffered (preferably pH>8) if necessary and the liquid diluent first rendered isotonic. These aqueous solutions are suitable for intravenous injection purposes. The oily solutions are suitable for intra-articular, intra-muscular and subcutaneous injection purposes. The preparation of all these solutions under sterile conditions is readily accomplished by standard pharmaceutical techniques well known to those skilled in the art. Additionally, it is also possible to administer the compounds of the present invention topically when treating inflammatory conditions of the skin and this may preferably be done by way of creams, jellies, gels, pastes, ointments and the like, in accordance with standard pharmaceutical practice.


## 15 Examples

The invention is illustrated in the following non-limiting examples in which, unless stated otherwise: all operations were carried out at room or ambient temperature, that is, in the range of 18-25 °C; evaporation of solvent was carried out using a rotary evaporator under reduced pressure with a bath temperature of up to 60 °C; reactions were monitored by thin layer chromatography (tlc) and reaction times are given for illustration only; melting points (m.p.) given are uncorrected (polymorphism may result in different melting points); the structure and purity of all isolated compounds were assured by at least one of the following techniques: tlc (Merck silica gel 60 F<sub>254</sub> precoated TLC plates or Merck NH<sub>2</sub> F<sub>254s</sub> precoated HPTLC plates), mass spectrometry, nuclear magnetic resonance (NMR), infrared red absorption spectra (IR), microanalysis or powder X-ray diffraction(PXRD) pattern. Yields are given for illustrative purposes only. Flash column chromatography was carried out using Merck silica gel 60 (230-400 mesh ASTM) or Fuji Silysia Chromatorex<sup>®</sup> DU3050 (Amino Type, 30-50 µm). Low-resolution mass spectral data (EI) were obtained on a Integrity (Waters) mass spectrometer or a Automass 120 (JEOL) mass spectrometer. Low-resolution mass spectral data (ESI) were obtained on a ZMD2 (Waters) mass spectrometer or a Quattro II (Micromass) mass


spectrometer. NMR data was determined at 270 MHz (JEOL JNM-LA 270 spectrometer) or 300 MHz (JEOL JNM-LA300) using deuterated chloroform (99.8% D) or dimethylsulfoxide (99.9% D) as solvent unless indicated otherwise, relative to tetramethylsilane (TMS) as internal standard in parts per million (ppm); conventional abbreviations used are: s = singlet, d = doublet, t = triplet, q = quartet, m = multiplet, br. = broad, etc. IR spectra were measured by a Shimadzu infrared spectrometer (IR-470). Optical rotations were measured using a JASCO DIP-370 Digital Polarimeter (Japan Spectroscopic CO, Ltd.). PXRD pattern was determined using a Rigaku RINT-TTR powder X-ray diffractometer fitted with an automatic sample changer, a 2 theta goniometer, beam divergence slits, a secondary monochromator and a scintillation counter. The sample was prepared for analysis by packing the powder on to an aluminum sample holder. The specimen was rotated by 60.00rpm and scanned by 4°/min. Chemical symbols have their usual meanings; b.p. (boiling point), m.p. (melting point), l (liter(s)), ml (milliliter(s)), g (gram(s)), mg(milligram(s)), mol (moles), mmol (millimoles), eq. (equivalent(s)).

**EXAMPLE 1:**

***N*-(1-[(4-Hydroxytetrahydro-2*H*-pyran-4-yl)methyl]piperidin-4-yl)methyl)-3-isopropyl-2-oxo-2,3-dihydro-1*H*-benzimidazole-1-carboxamide hydrochloride**



**Step 1. *tert*-Butyl (1-[(4-hydroxytetrahydro-2*H*-pyran-4-yl)methyl]piperidin-4-yl)methyl)carbamate**



To a stirred solution of *tert*-butyl (piperidin-4-ylmethyl)carbamate (22.3 g, 104 mmol) in methanol was added 1,6-dioxaspiro[2.5]octane (14.2 g, 124 mmol, Satyamurthy, Nagichettiar et al., *Phosphorus Sulfur*, 1984, 19, 113) at ambient temperature.

Then, the mixture was heated at 60°C for 4h. The volatile components were

removed by evaporation and the resulting viscous oil was precipitated with a mixture of hexane and diethylether. The precipitate was collected by filtration and recrystallized with a mixture of hexane and 2-propanol to give title compound 14.2 g (42%) as a colorless powder.

5 MS (ESI)  $m/z$ : 329 ( $M+H^+$ ).


m.p.: 104°C.

<sup>1</sup>H-NMR ( $CDCl_3$ )  $\delta$ : 1.23-1.31 (2 H, m), 1.44 (9 H, s), 1.51-1.69 (8 H, m), 2.27-2.38 (4 H, m), 2.83-2.88 (2 H, m), 3.00 (2 H, t,  $J=6.2$  Hz), 3.70-3.85 (4 H, m).

Anal. Calcd. for  $C_{17}H_{32}N_2O_4$ : C, 62.17; H, 9.82; N, 8.53. Found: C, 62.07; H, 9.92; N,

10 8.58.

Step 2. 4-({[4-(Aminomethyl)piperidin-1-yl]methyl}tetrahydro-2H-pyran-4-ol



To a solution of *tert*-butyl ({1-[(4-hydroxytetrahydro-2H-pyran-4-yl)methyl]piperidin-4-yl)methyl}carbamate (50.28 g, 153 mmol) in methanol was

15 added 4N HCl in dioxane (200 mL, 800 mmol) at room temperature. After 4h, the volatile materials were removed by evaporation. The resulting amorphous was

precipitated with diethyl ether/methanol (5:1). The precipitate was collected and added to the ice cooled 6N NaOH aq. (200 mL) gradually. The mixture was extracted with dichloromethane/methanol (10:1) for 4 times. The combined organic phase was


20 washed with brine, dried over  $MgSO_4$  and concentrated to give 24.90 g (99%) of the title compound as pale brown amorphous.

MS (ESI)  $m/z$ : 229 ( $M+H^+$ ).

<sup>1</sup>H-NMR ( $CDCl_3$ )  $\delta$ : 1.19-1.28 (2 H, m), 1.44-1.63 (8 H, m), 1.65-1.71 (2 H, m), 2.32 (2 H, s), 2.35 (2 H, t,  $J=11.0$  Hz), 2.57 (2 H, d,  $J=5.7$  Hz), 2.85-2.90 (2 H, m), 3.70-

25 3.81 (4 H, m).

Step 3. *N*-({1-[(4-Hydroxytetrahydro-2H-pyran-4-yl)methyl]piperidin-4-yl)methyl}-3-isopropyl-2-oxo-2,3-dihydro-1H-benzimidazole-1-carboxamide





- To a stirred mixture of 1-isopropyl-1,3-dihydro-2H-benzimidazol-2-one (*J. Med. Chem.* **1999**, 42, 2870-2880) (23.0 g, 130 mmol) and triethylamine (54.6 mL, 392 mmol) in tetrahydrofuran (300 mL) was added triphosgen (38.8 g, 130 mmol) in tetrahydrofuran (200 mL) gradually at room temperature. Then, the mixture was
- 5 heated at 80°C for 4h. After cooling, a solution of 4-([4-(aminomethyl)piperidin-1-yl]methyl)tetrahydro-2H-pyran-4-ol (step 2 of Example 1) (24.9 g, 109 mmol) and triethylamine (45 mL, 109 mmol) in tetrahydrofuran (500 mL) was added to the mixture. Then, the mixture was heated at 80°C for 6h. After cooling, sat. NaHCO<sub>3</sub> aq was added to the mixture. The mixture was extracted with ethyl acetate (500 mLx4).
- 10 The extracts were washed with brine, dried over MgSO<sub>4</sub> and concentrated. The residue was chromatographed on a column of aminopropyl-silica gel eluting with hexane/ethyl acetate (3:1) to give 31.3 g (67%) of the title compound as a white solid.
- <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 8.80 (1 H, br t, *J* = 6.0 Hz), 8.06 (1 H, m), 7.41 (1 H, m), 7.19 (1 H, dt, *J* = 1.5, 7.7 Hz), 7.12 (1 H, dt, *J* = 1.3, 7.7 Hz), 4.64 (1 H, septet, *J* = 7.0 Hz),
- 15 4.08 (1 H, br s), 3.68-3.44 (4 H, m), 3.19 (2 H, t, *J* = 6.0 Hz), 2.89 (2 H, m), 2.20 (2 H, br s), 2.09 (2 H, m), 1.68-1.10 (9 H, m), 1.47 (6 H, d, *J* = 7.0 Hz).
- MS (ESI) *m/z*: 431 (M+H<sup>+</sup>).
- Anal. Calcd. for C<sub>23</sub>H<sub>34</sub>N<sub>4</sub>O<sub>4</sub>: C, 64.16; H, 7.96; N, 13.01. Found: C, 64.13; H, 7.97; N, 12.99.

- 20 Step 4. *N*-({1-[(4-Hydroxytetrahydro-2H-pyran-4-yl)methyl]piperidin-4-yl}methyl)-3-isopropyl-2-oxo-2,3-dihydro-1H-benzimidazole-1-carboxamide hydrochloride


- To a stirred solution of *N*-({1-[(4-hydroxytetrahydro-2H-pyran-4-yl)methyl]piperidin-4-yl}methyl)-3-isopropyl-2-oxo-2,3-dihydro-1H-benzimidazole-1-carboxamide (27.0 g, 62 mmol) in methanol (150 mL) was added 10% HCl-
- 25 methanol (100 mL) at ambient temperature. After 30min, the volatile materials were removed by evaporation. The resulting amorphous was precipitated by ethanol/diethylether. The precipitate was recrystallized from ethanol/diethylether (1:1) to give 26.5 g (90 %) of title compound as a colorless powder.
- <sup>1</sup>H-NMR (DMSO-*d*<sub>6</sub>) δ: 1.49 (6 H, d, *J*=6.9 Hz), 1.50-1.70 (4 H, m), 1.76-1.91 (5 H, m), 3.00-3.12 (3 H, m), 3.15-3.45 (3 H, m), 3.60-3.70 (6 H, m), 4.61-4.69 (1 H, m),
- 30 5.46-5.49 (1 H, m), 7.13 (1 H, t, *J*=7.8 Hz) 7.20 (1 H, t, *J*=7.8 Hz), 7.42 (1 H, d, *J*=7.9 Hz), 8.07 (1 H, d, *J*=8.0 Hz), 8.86 (1 H, m), 9.61-9.81 (1 H, m)

MS (ESI)  $m/z$ : 431 ( $M+H^+$ ).

Anal. Calcd. for  $C_{23}H_{35}N_4O_4Cl$ : C, 59.15; H, 7.55; N, 2.00. Found: C, 58.81; H, 7.57; N, 11.85.


- Alternative route to synthesize 4-{{[4-(aminomethyl)piperidin-1-yl]methyl}tetrahydro-2H-pyran-4-ol} is described below.

Step 1. Benzyl ({1-[(4-hydroxytetrahydro-2H-pyran-4-yl)methyl]piperidin-4-yl)methyl}carbamate



- A mixture of benzyl (piperidin-4-ylmethyl)carbamate (7.77 g, 31.3 mmol, Bose, D. Subhas et al, *Tetrahedron Lett.*, 1990, 31, 6903) and 1,6-dioxaspiro[2.5]octane (4.29 g, 37.6 mmol, Satyamurthy, Nagichettiar et al., *Phosphorus Sulfur*, 1984, 19, 113) in methanol (93 mL) was stirred at room temperature for 20 h. Then the mixture was refluxed for 8 h. After cooling to room temperature, the solvent was removed *in vacuo*. The residue was chromatographed on a column of silica gel eluting with methanol/dichloromethane (1:20) to give 5.60 g (49%) of the title compound as a colorless oil.

Step 2. 4-{{[4-(Aminomethyl)piperidin-1-yl]methyl}tetrahydro-2H-pyran-4-ol}



- A mixture of benzyl ({1-[(4-hydroxytetrahydro-2H-pyran-4-yl)methyl]piperidin-4-yl)methyl}carbamate (5.60 g, 15.5 mmol, step1) and palladium on activated carbon (10 wt.%, 1.20 g) in methanol (250 mL) was hydrogenated at room temperature for 20 h. Then, the mixture was filtered through a pad of Celite, and the filtrate was concentrated *in vacuo* to give 3.30 g (94%) of the title compound as slightly yellow oil.

- Following is another route to synthesize 4-{{[4-(aminomethyl)piperidin-1-yl]methyl}tetrahydro-2H-pyran-4-ol}.

Step 1. 1-[(4-hydroxytetrahydro-2H-pyran-4-yl)methyl]piperidine-4-carboxamide

The mixture of trimethylsulfoxonium iodide (0.791 g, 3.52 mmol) and 2N-NaOH aq (1.76 mL, 3.52 mmol) in acetonitrile (1.62 mL) was stirred at 50 °C for 30 min. Then to the mixture was added tetrahydro-4H-pyran-4-one (0.324 g, 3.20 mmol) and the resulting mixture was stirred at 50°C for 3 h. Sat. NaCl aq. (10 mL) was added to the reaction mixture at room temperature and organic layer was extracted with CH<sub>2</sub>Cl<sub>2</sub> (20 mL), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated.

After removal of the solvent, MeOH (1.62 mL) and isonipecotamide (0.381 g, 2.88 mmol) were added to the residue, the mixture was stirred at 75°C for 14 h under N<sub>2</sub>.

The reaction mixture was concentrated and the residue was recrystallized from MeOH-acetonitrile to give 0.484 g (2.00 mmol) of title compound as a white solid.

<sup>1</sup>H-NMR (300 MHz, DMSO-*d*<sub>6</sub>) δ 7.19 (br s, 1H), 6.69 (br s, 1 H), 4.10 (s, 1 H), 3.70-3.50 (m, 4 H), 2.95-2.85 (m, 2 H), 2.20 (s, 2 H), 2.15-1.85 (m, 3 H), 1.65-1.50 (m, 6 H), 1.40-1.25 (m, 2 H).

Step 2. 4-[[4-(Aminomethyl)piperidin-1-yl]methyl]tetrahydro-2H-pyran-4-ol tosylate

To a stirred suspension of NaBH<sub>4</sub> (0.505 g, 13.2 g) in triethylene glycol dimethyl ether (12.8 mL) was added the solution 1-[(4-hydroxytetrahydro-2H-pyran-4-yl)methyl]piperidine-4-carboxamide (0.640 g, 2.64 mmol) and AcOH (0.765 mL, 13.2 mmol) in triethylene glycol dimethyl ether (3.2 mL) dropwise at 80°C under N<sub>2</sub>.

The reaction mixture was quenched with 2N-HCl aq until pH value was < 3, then the resulting mixture was stirred at room temperature for 1 h. To the mixture CH<sub>2</sub>Cl<sub>2</sub> (30 mL) and 2N-NaOH aq. was added until pH value of aqueous layer was > 10. Organic layer was extracted with CH<sub>2</sub>Cl<sub>2</sub> for three times, and the combined organic layer was dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated.


To the residual solution (title compound in triethylene glycol dimethyl ether) the solution of *p*-toluenesulfonic acid monohydrate (0.408 g, 2.11 mmol) in MeOH (1.28 mL) was added at 60°C, then the mixture was cooled to room temperature. Appeared solids were collected by suction and wash with hexane to give title compound (0.340 g, 0.849 mmol) as a white solid

<sup>1</sup>H-NMR (300 MHz, DMSO-*d*<sub>6</sub>) δ 7.61 (br s, 2 H), 7.55-7.40 (m, 2 H), 7.15 -7.05 (m, 2 H), 4.11 (br s, 1 H), 3.70-3.45 (m, 4 H), 2.95-2.85 (m, 2 H), 2.68 (d, J = 7.0, 2 H), 2.29 (s, 3 H), 2.22 (s, 2 H), 2.07 (t, J = 11.0, 2 H), 1.65-1.45 (m, 4 H), 1.55-1.35 (m, 1

H), 1.40-1.25 (m, 2 H), 1.30-1.10 (m, 2 H).

# EXAMPLE 2.

*N*-({1-[(4-Hydroxytetrahydro-2*H*-pyran-4-yl)methyl]piperidin-4-yl)methyl}-3-isopropyl-2-oxo-2,3-dihydro-1*H*-benzimidazole-1-carboxamide hemiedisylate



To a stirred solution of *N*-({1-[(4-hydroxytetrahydro-2*H*-pyran-4-yl)methyl]piperidin-4-yl)methyl}-3-isopropyl-2-oxo-2,3-dihydro-1*H*-benzimidazole-1-carboxamide 1.51 g (3.51 mmol) in ethyl acetate (10 mL) and methanol (10 mL) was added a solution of 1,2-ethanedithiolic acid dihydrate 397 mg (1.75 mmol) in methanol (5.0 mL) and the resulting suspension was stirred for 5 h at room temperature. The mixture was filtered and the first crop was dried under vacuum for 5 h at 100 °C to give 1.78 g of crude product. 1.61 g of the crude product was dissolved in methanol (20 mL) and ethyl acetate (20 mL) was added to the solution. The resulting suspension was stirred for 2 h at room temperature. The mixture was filtered and the crop was dried under vacuum for 4 h at 100 °C to give the titled compound 1.13 g (61%) as colorless crystals.

MS (ESI) *m/z*: 431 (M+H)<sup>+</sup>.

*m.p.*: 233 °C.


IR (KBr) *v*: 2866, 1738, 1683, 1558, 1373, 1217, 1028, 756 cm<sup>-1</sup>.

<sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 8.96 (0.25 H, br s), 8.85 (1 H, br t, *J* = 6.0 Hz), 8.61 (0.75 H, br s), 8.06 (1 H, m), 7.43 (1 H, m), 7.21 (1 H, dt, *J* = 1.3, 7.7 Hz), 7.13 (1 H, dt, *J* = 1.2, 7.7 Hz), 5.26 (1 H, br s), 4.65 (1 H, septet, *J* = 7.0 Hz), 3.74-2.92 (12 H, m), 2.64 (2 H, s), 2.00-1.35 (9 H, m), 1.47 (6 H, d, *J* = 7.0 Hz).

Anal. calcd. for C<sub>23</sub>H<sub>34</sub>N<sub>4</sub>O<sub>4</sub> • 0.5 C<sub>2</sub>H<sub>6</sub>O<sub>6</sub>S<sub>2</sub>: C, 54.84; H, 7.09; N, 10.66; S, 6.10.

Found: C, 54.50; H, 7.24; N, 10.60; S, 6.08.

PXRD pattern angle (2-Theta°): 10.2, 11.9, 16.3, 17.3, 17.6, 21.8, 24.2.

**EXAMPLE 3.****3-Isopropyl-N-[[1-(2-morpholin-4-yl-2-oxoethyl)piperidin-4-yl]methyl]-2-oxo-2,3-dihydro-1H-benzimidazole-1-carboxamide monooxalate**

The tilted compound was prepared with the similar method shown in the Step 3 of Preparation 1 by using 4-(chloroacetyl)morpholine (B. G. Hazra; V. S. Pore; S. P. Maybhat, *Org. Prep. Proced. Int.*, **1989**, 21, 355-8).

MS (ESI)  $m/z$ : 440 (M+H)<sup>+</sup>.


10 m.p.: 194.2 °C.

IR (KBr)  $\nu$ : 3443, 2934, 1765, 1728, 1686, 1659, 1612, 1551 cm<sup>-1</sup>.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) (free base)  $\delta$ : 9.00-8.88 (1H, m) 8.30-8.22 (1H, m), 7.23-7.12 (3H, m), 4.78-4.62 (1H, m), 3.66 (4H, s), 3.70-3.58 (4H, m), 3.32 (2H, t, J=6.3 Hz), 3.15 (2H, s), 2.94-2.84 (2H, m), 2.14-2.01 (2H, m), 1.86-1.23 (5H, m), 1.56 (6H, d, J=7.0 Hz).

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) (salt form)  $\delta$ : 8.92-8.80 (1H, m) 8.07 (1H, d, J=7.7 Hz), 7.45 (1H, d, J=7.5 Hz), 7.26-7.06 (2H, m), 4.76-4.56 (1H, m), 4.10-2.60 (18H, m), 1.90-1.40 (3H, m), 1.49 (6H, d, J=6.9 Hz).


Anal. Calcd. for C<sub>25</sub>H<sub>33</sub>N<sub>5</sub>O<sub>8</sub>: C, 56.27; H, 6.61; N, 13.13. Found: C, 56.25; H, 6.82; N, 12.98.

**EXAMPLE 4.****3-Isopropyl-N-[[1-(3-morpholin-4-yl-3-oxopropyl)piperidin-4-yl]methyl]-2-oxo-2,3-dihydro-1H-benzimidazole-1-carboxamide monooxalate**

- A mixture of 3-isopropyl-2-oxo-*N*-(piperidin-4-ylmethyl)-2,3-dihydro-1*H*-benzimidazole-1-carboxamide (150 mg, 0.474 mmol) and 4-(3-chloro-propanoyl)-morpholine (G. Mattalia; C. Serafini; U. Bucciarelli, *Farmaco, Ed. Sci.*, **1976**, *31*, 457-67) (300 mg, 1.185 mmol) in 4.7 ml *N,N*-dimethylformamide was added
- 5 triethylamine (0.23 ml, 1.659 mmol) and sodium iodide (178 ml, 1.185 mmol). The reaction mixture was stirred at 90 °C for 6 days. The reaction mixture was then concentrated by evaporation. The residue was diluted aqueous NaHCO<sub>3</sub> 10 ml, extracted with dichloromethane 30 ml for three times. The combined extract was dried over MgSO<sub>4</sub> and concentrated. Preparative TLC (eluent:
- 10 CH<sub>2</sub>Cl<sub>2</sub>/methanol=10/1) afforded a brown amorphous oil 130 mg (60%). The amorphous (130 mg) was dissolved in 3 ml methanol and acidified with a solution of 24 mg oxalic acid in 2 ml MeOH. The mixture was concentrated. Crystallization of the resulting residue with AcOEt-EtOH afforded a white amorphous 107 mg as the titled compound.
- 15 MS (ESI) *m/z*: 458 (M+H)<sup>+</sup>.  
 IR (KBr)  $\nu$ : 3443, 2941, 1732, 1697, 1686, 1647, 1638, 1558 cm<sup>-1</sup>.  
<sup>1</sup>H-NMR (CDCl<sub>3</sub>) (free base)  $\delta$ : 9.06-8.94 (1H, br) 8.24-8.19 (1H, m), 7.26-7.10 (3H, m), 4.76-4.64 (1H, m), 3.75-2.80 (10H, m), 2.60-1.30 (13H, m), 1.56 (6H, d, J=7.0 Hz).
- 20 <sup>1</sup>H-NMR (CDCl<sub>3</sub>) (salt form)  $\delta$ : 9.10-9.00 (1H, m) 8.27-8.17 (1H, m), 7.33-7.12 (3H, m), 4.87-4.62 (1H, m), 3.78-2.65 (16H, m), 2.20-1.60 (7H, m), 1.56 (6H, d, J=6.9 Hz).  
 Anal. Calcd. for C<sub>26</sub>H<sub>37</sub>N<sub>5</sub>O<sub>8</sub>•0.9C<sub>2</sub>H<sub>2</sub>O<sub>4</sub>•1.3H<sub>2</sub>O: C, 51.21; H, 6.40; N, 10.74.  
 Found: C, 50.90; H, 6.26; N, 11.13.

## 25 EXAMPLE 5.

**3-Isopropyl-*N*-{[1-(4-morpholin-4-yl-4-oxobutyl)piperidin-4-yl]methyl}-2-oxo-2,3-dihydro-1*H*-benzimidazole-1-carboxamide monooxalate**



The tilted compound was prepared with the similar method shown in the example 4 by using 4-(4-chloro-butyl)-morpholine (Schlesinger; Prill; B. G. Hazra; *J.Amer.Chem.Soc.*, **1956**, *78*, 6123-6124).

MS (ESI)  $m/z$ : 472 ( $M+H$ )<sup>+</sup>.

- 5 IR (KBr)  $\nu$ : 3443, 1728, 1686, 1647-1616, 1551  $\text{cm}^{-1}$ .


<sup>1</sup>H-NMR ( $\text{CDCl}_3$ ) (free base)  $\delta$ : 9.02-8.88 (1H, m), 8.31-8.20 (1H, m), 7.22-7.04 (3H, m), 4.80-4.60 (1H, m), 3.66-3.56 (8H, m), 3.40-3.22 (2H, m), 3.00-2.88 (2H, m), 2.50-2.30 (6H, m), 2.00-1.20 (7H, m), 1.57 (6H, d,  $J=7.1$  Hz).

- <sup>1</sup>H-NMR ( $\text{DMSO}-d_6$ ) (salt form)  $\delta$ : 8.93-8.79 (1H, m), 8.07 (1H, d,  $J=7.5$  Hz), 7.44 (1H, d,  $J=7.5$  Hz), 7.27-7.08 (2H, m), 4.75-4.58 (1H, m), 4.47-2.30 (18H, m), 1.90-0.90 (7H, m), 1.49 (6H, d,  $J=6.9$  Hz).


Anal. Calcd. for  $\text{C}_{27}\text{H}_{39}\text{N}_5\text{O}_8$ : C, 57.74; H, 7.00; N, 12.47. Found: C, 57.52; H, 7.03; N, 12.32.

# 15 EXAMPLE 6.

*N*-{1-[(*trans*-1,4-Dihydroxyhexyl)methyl]piperidin-4-yl)methyl}-3-isopropyl-2-oxo-2,3-dihydro-1*H*-benzimidazole-1-carboxamide hydrochloride



Step 1. *tert*-Butyl(1-oxaspiro[2.5]oct-6-yloxy)diphenylsilane



- 20 To a stirred suspension of sodium hydride (60% in mineral oil, 441 mg, 11.0 mmol) in DMSO (7 ml) was added trimethylsulfoxonium iodide (2.53 g, 11.5 mmol) at room temperature, and the mixture was stirred at room temperature for 30min. To this mixture was added a solution of 4-[[*tert*-butyl(diphenyl)silyl]oxy]cyclohexanone (Okamura, William H. et al., *J.Org.Chem.*, **1993**, *58*, 600-610, 3.53 g, 10.0 mmol) in
- 25 DMSO (35 ml) dropwise at room temperature, the mixture was stirred at room

- temperature for 2 h. Then the mixture was diluted with water (600 ml), and extracted with diethylether (200 ml x 4). The combined organic layer was dried over magnesium sulfate, and concentrated *in vacuo*. The residue was chromatographed on a column of silica gel eluting with *n*-hexane/ethyl acetate (1:10), and then purified with PTLC eluting with *n*-hexane/ethyl acetate (1:15) to give 459 mg (13%, *trans*) and 390 mg (11%, *cis*) of the title compound as colorless oil respectively.


(*trans*)

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 7.70-7.66 (4 H, m), 7.46-7.35 (6 H, m), 4.03-3.97 (1 H, m), 2.63 (2 H, s), 2.07-1.63 (8 H, m), 1.08 (9 H, s).

- 10 (*cis*)

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 7.70-7.65 (4 H, m), 7.46-7.35 (6 H, m), 3.97-3.83 (1 H, m), 2.58 (2 H, s), 1.83-1.37 (8 H, m), 1.07 (9 H, s).

Step 2. *N*-{[1-({*trans*-4-[*tert*-Butyl(diphenyl)silyl]oxy-1-hydroxycyclohexyl)methyl]piperidin-4-yl)methyl}-3-isopropyl-2-oxo-2,3-dihydro-1*H*-benzimidazole-1-carboxamide.




- A mixture of *tert*-butyl[(3*R*,6*R*)-1-oxaspiro[2.5]oct-6-yloxy]diphenylsilane (Step1, *trans*-isomer, 283.0 mg, 0.772mmol) and 3-isopropyl-2-oxo-*N*-(piperidin-4-ylmethyl)-2,3-dihydro-1*H*-benzimidazole-1-carboxamide (Preparation 1, step 2, 2.48g, 0.0194mol) in MeOH (4 ml) was heated at 50°C with stirring for 2 days. After cooling, the reaction mixture was evaporated to remove the solvent, and residue was chromatographed on a column of silica gel eluting with ethyl acetate/*n*-hexane (1:10) then methanol/dichloromethane (1:20) to give 308.1 mg (58 %) of the title compound as a colorless syrup.
- 20 MS (ESI) *m/z*: 683 (M+H)<sup>+</sup>.
- <sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 8.93 (1 H, m), 8.32-8.23 (1 H, m), 7.72-7.60 (4 H, m), 7.46-7.32 (6 H, m), 7.22-7.10 (3 H, m), 4.80-4.62 (1 H, m), 3.96 (1 H, m), 3.31 (2 H, t, *J*=6.26Hz), 2.92 (2 H, d, *J*=10.88 Hz), 2.45-2.29 (4 H, m), 1.85-1.65 (6 H, m), 1.65-1.43 (9 H, m, including 6 H, d, *J*=7.09 Hz at 1.56 ppm), 1.43-1.25 (4 H, m), 1.06 (9 H,



s).

step 3. N-([1-([trans-1,4-Dihydroxycyclohexyl)methyl]piperidin-4-yl)methyl]-3-isopropyl-2-oxo-2,3-dihydro-1H-benzimidazole-1-carboxamide hydrochloride



- 5 A mixture of *tert*-butyl N-([1-([trans-4-[*tert*-butyl(diphenyl)silyl]oxy-1-hydroxycyclohexyl)methyl]piperidin-4-yl)methyl]-3-isopropyl-2-oxo-2,3-dihydro-1H-benzimidazole-1-carboxamide (234 mg, 0.343mol) and HCl solution of MeOH (50ml) was stirred at room temperature for 4 h. Then the solvent was removed *in vacuo*. The residue was basified with saturated aqueous NaHCO<sub>3</sub> (30ml), extracted with CH<sub>2</sub>Cl<sub>2</sub> (30 ml×3 times) and the combined organic layer was dried over Na<sub>2</sub>SO<sub>4</sub>. Removal of the solvent gave a residue, which was chromatographed on a column of NH-silica gel eluting with ethyl acetate /*n*-hexane (1:1-2:1) to give 140.1 mg (92%) of the title compound as a colorless syrup.

MS (ESI) *m/z*: 445 (M+H)<sup>+</sup>.

- 15 <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 8.93 (1 H, br t, J=5.87 Hz), 8.32-8.20 (1 H, m), 7.25-7.03 (3 H, m), 4.80-4.62 (1 H, m), 3.94 (1 H, m), 3.31 (2 H, t, J=6.10 Hz), 2.89 (2 H, br d, J=11.53 Hz), 2.36 (2 H, s), 2.34 (2 H, t, J=11.86 Hz), 2.00-1.85 (2 H, m), 1.82-1.25 (18 H, m, including 6 H, d, J=7.09 Hz at 1.56 ppm).

- 140.1 mg of this syrup was dissolved in HCl solution in MeOH (4 ml), concentrated, and dried *in vacuo* at 50°C for 5h to give 139.2 mg of title compound as a yellow amorphous solid.

MS (ESI) *m/z*: 445 (M+H)<sup>+</sup>.

- <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ : 9.35-8.75 (1H, m), 8.86 (1 H, t, J=6.59 Hz), 8.07 (1 H, d, J=7.74 Hz), 7.44 (1 H, d, J=7.58 Hz), 7.22 (1 H, dt, J=1.15 Hz, 7.42 Hz), 7.14 (1 H, dt, J=1.32 Hz, 7.74 Hz), 5.04 (1 H, br s), 4.75-4.45 (1 H, m), 3.70 (1 H, br s), 3.59 (2 H, d, J=11.70 Hz), 3.50-2.90 (8 H, m), 1.90-1.57 (8 H, m), 1.57-1.30 (10 H, m, including 6 H, d, J=6.92 Hz at 1.49 ppm)


IR(KBr): 3285, 2936, 2677, 1728, 1686, 1611, 1549, 1481, 1375, 1298, 1204, 1157, 1101, 1018, 762 cm<sup>-1</sup>

Anal. Calcd for  $C_{24}H_{36}N_4O_4 \cdot HCl \cdot 2H_2O$ : C, 57.76; H, 7.88; N, 11.23. Found: C, 57.54; H, 7.90; N, 11.21.

PXRD pattern angle (2-Theta°): 8.3, 14.5, 17.7, 18.3, 19.1, 26.4, 27.5.


## 5 EXAMPLE 7.

*N*-({1-[(*cis*-1,4-Dihydroxyhexyl)methyl]piperidin-4-yl)methyl}-3-isopropyl-2-oxo-2,3-dihydro-1*H*-benzimidazole-1-carboxamide hydrochloride



Step 1. *N*-{[1-({*cis*-4-[*tert*-Butyl(diphenyl)silyl]oxy-1-

- 10 hydroxycyclohexyl)methyl]piperidin-4-yl)methyl}-3-isopropyl-2-oxo-2,3-dihydro-1*H*-benzimidazole-1-carboxamide.




The title compound was prepared according to the procedure described of Step 2 in the Example 6 using *tert*-butyl[(3*S*,6*S*)-1-oxaspiro[2.5]oct-6-

- 15 yloxy]diphenylsilane (Example 6, Step 1, *cis*-isomer, 311.0 mg, 0.848 mmol) instead of *tert*-butyl[(3*R*,6*R*)-1-oxaspiro[2.5]oct-6-yloxy]diphenylsilane.

MS (ESI) *m/z*: 683 (*M*+*H*)<sup>+</sup>.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 8.91 (1 H, t, *J*=5.87 Hz), 8.30-8.22 (1 H, m), 7.72-7.63 (4 H, m), 7.45-7.30 (6 H, m), 7.20-7.10 (3 H, m), 4.80-4.63 (1 H, m), 3.59 (1 H, m), 3.29 (2 H, t, *J*=6.24 Hz), 2.83 (2 H, d, *J*=11.74 Hz), 2.26 (2 H, t, *J*=11.55 Hz), 2.18 (2 H, s), 1.85-1.65 (4 H, m), 1.65-1.50 (11 H, m, including 6 H, d, *J*=7.15 Hz at 1.56 ppm), 1.40-1.30 (2 H, m), 1.15-1.00 (11 H, m, including 9 H, s, 1.05 ppm).

step 2. *N*-{[1-[(*cis*-1,4-Dihydroxycyclohexyl)methyl]piperidin-4-yl)methyl}-3-isopropyl-2-oxo-2,3-dihydro-1*H*-benzimidazole-1-carboxamide hydrochloride



The title compound was prepared according to the procedure described of Step 3 in the Example 6 using *N*-{[1-(*cis*-4-[*tert*-butyl(diphenyl)silyl]oxy-1-hydroxycyclohexyl)methyl]piperidin-4-yl)methyl}-3-isopropyl-2-oxo-2,3-dihydro-1*H*-benzimidazole-1-carboxamide (295.0 mg, 0.432 mmol) instead of *N*-{[1-(*trans*-4-[diphenyl(trimethylsilyl)methoxy]-1-hydroxycyclohexyl)methyl]piperidin-4-yl)methyl}-3-isopropyl-2-oxo-2,3-dihydro-1*H*-benzimidazole-1-carboxamide. MS (ESI) *m/z*: 445 (*M*+*H*)<sup>+</sup>.

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 8.93 (1 H, br t, *J*=5.60 Hz), 8.31-8.22 (1 H, m), 7.25-7.10 (3 H, m), 4.80-4.62 (1 H, m), 3.63-3.49 (1 H, m), 3.31 (2 H, t, *J*=6.10 Hz), 2.89 (2 H, br d, *J*=11.54 Hz), 2.33 (2 H, dt, *J*=1.81 Hz, 11.70 Hz), 1.85-1.60 (16 H, m, including 6 H, d, *J*=7.09 Hz at 1.57 ppm), 1.45-1.18 (4 H, m).

165.7 mg of this syrup was dissolved in HCl solution in MeOH (4 ml), concentrated, and dried *in vacuo* at 50°C for 5h to give 164.7 mg of title compound as a yellow amorphous solid.

MS (ESI) *m/z*: 445 (*M*+*H*)<sup>+</sup>.


<sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ : 9.30-8.90 (1H, m), 8.86 (1 H, t, *J*=5.93 Hz), 8.07 (1 H, d, *J*=7.58 Hz), 7.44 (1 H, d, *J*=7.58 Hz), 7.22 (1 H, dt, *J*=1.48 Hz, 7.75 Hz), 7.15 (1 H, dt, *J*=1.15 Hz, 7.74 Hz), 4.75-4.58 (1 H, m), 3.70-2.90 (11 H, m), 1.90-1.67 (6 H, m), 1.67-1.20 (12 H, m, including 6 H, d, *J*=6.92 Hz at 1.49 ppm).

IR(KBr): 3294, 2936, 2673, 1728, 1686, 1611, 1545, 1479, 1375, 1298, 1203, 1158, 1134, 1101, 1051, 762 cm<sup>-1</sup>


Anal. Calcd for C<sub>24</sub>H<sub>36</sub>N<sub>4</sub>O<sub>4</sub>·HCl·5H<sub>2</sub>O: C, 54.79; H, 8.05; N, 10.65. Found: C, 54.75; H, 7.88; N, 10.56.

#### EXAMPLE 8:

**6-Fluoro-*N*-{[1-(4-(4-hydroxytetrahydro-2*H*-pyran-4-yl)methyl]piperidin-4-yl)methyl}-3-isopropyl-2-oxo-2,3-dihydro-1*H*-benzimidazole-1-carboxamide hydrochloride**



Step 1. 6-Fluoro-N-({1-[(4-hydroxytetrahydro-2H-pyran-4-yl)methyl]piperidin-4-yl)methyl}-3-isopropyl-2-oxo-2,3-dihydro-1H-benzimidazole-1-carboxamide



- 5 The title compound was prepared according to the procedure described in Step 3 of Example 1 from 5-fluoro-1-isopropyl-1,3-dihydro-2H-benzimidazol-2-one [Tapia et al., *J. Med. Chem.*, 1999, 42, 2880.] and 4-[[4-(aminomethyl)piperidin-1-yl]methyl]tetrahydro-2H-pyran-4-ol (step 2 of Example 1).  
MS (ESI) m/z: 449 (M+H<sup>+</sup>).

- 10 <sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.12-1.70 (8 H, m), 1.55 (6 H, d, J=7.0 Hz), 1.74 (2 H, brd, 12.8 Hz), 2.31 (2 H, s), 2.35 (2 H, brt, J=11.9 Hz), 2.88 (2 H, brd, J=11.7 Hz), 3.30 (2 H, t, J=6.2 Hz), 3.70-3.85 (4 H, m), 4.62-4.75 (1 H, m), 6.90 (1 H, td, J=9.0, 2.4 Hz), 7.02-7.07 (1 H, m), 8.05 (1H, dd, J=9.5, 2.6 Hz), 8.85-8.92 (1 H, m).

- 15 Step 2. 6-Fluoro-N-({1-[(4-hydroxytetrahydro-2H-pyran-4-yl)methyl]piperidin-4-yl)methyl}-3-isopropyl-2-oxo-2,3-dihydro-1H-benzimidazole-1-carboxamide hydrochloride

- The title compound was prepared according to the procedure described in Step 4 of Example 1 from 6-fluoro-N-({1-[(4-hydroxytetrahydro-2H-pyran-4-yl)methyl]piperidin-4-yl)methyl}-3-isopropyl-2-oxo-2,3-dihydro-1H-benzimidazole-1-carboxamide (step 1 of Example 8).

- MS (ESI) m/z: 449 (M+H<sup>+</sup>).  
<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 1.46 (6 H, d, J=6.9 Hz), 1.55-1.65 (4 H, m), 1.70-1.91 (4 H, m), 2.90-3.28 (8 H, m), 3.50-3.67 (6 H, m), 4.56-4.69 (1 H, m), 5.30-5.37 (1 H, m), 5.76 (1 H, s), 7.08 (1 H, td, J=9.0, 2.4 Hz), 7.44-7.49 (1 H, m), 7.85 (1H, dd, J=9.5, 2.5 Hz), 8.81-8.85 (1 H, m).


Anal. Calcd. for C<sub>23</sub>H<sub>34</sub>FN<sub>4</sub>O<sub>4</sub>Cl: C, 56.96; H, 7.07; N, 11.55. Found: C, 57.00; H, 7.20; N, 11.43.

PXRD pattern angle (2-Theta°): 10.0, 14.6, 16.2, 18.5, 23.2, 25.3, 27.3.

### EXAMPLE 9:

#### 5-fluoro-N-({1-[(4-hydroxytetrahydro-2H-pyran-4-yl)methyl]piperidin-4-

#### yl)methyl}-3-isopropyl-2-oxo-2,3-dihydro-1H-benzimidazole-1-carboxamide



#### Step 1. (5-fluoro-2-nitrophenyl)isopropylamine

To a stirred mixture of 2,4-difluoro-1-nitrobenzene (4.77 g, 30 mmol) and K<sub>2</sub>CO<sub>3</sub> (4.14 g, 30 mmol) in THF (30 mL) was added isopropyl amine (1.77g, 30 mmol) in THF (10 mL) at 0°C. After being stirred for 13h, the insoluble materials were removed by pad of Celite and the filtrate was concentrated under reduced pressure to give title compound (5.25 g, 88%) as a pale yellow oil.

MS (ESI) m/z: 405 (M+H<sup>+</sup>).

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 8.21 (1 H, dd, J= 9.3, 6.0 Hz), 6.48 (1 H, dd, J= 11.7, 2.6 Hz), 6.39-6.29 (1 H, m), 3.81-3.66 (1 H, m), 1.33 (6 H, d, J=6.4 Hz)

#### Step 2. 6-fluoro-1-isopropyl-1,3-dihydro-2H-benzimidazol-2-one

A mixture of (5-fluoro-2-nitrophenyl)isopropylamine (Step 1 of Example 9, 5.85 g, 30 mmol) and 10% Pd-C (600 mg) in MeOH was stirred under atmosphere of hydrogen gas at room temperature for 12 h. The catalyst was filtered off on a pad of Celite, and the filtrate was evaporated under reduced pressure. To the residue was added 1, 1'-carbonyldiimidazole (4.5 g, 28 mmol) and THF (100 mL) and then stirred at 100°C for 10h. After cooling, the volatile materials were removed under reduced pressure and the residue was partitioned between ethylacetate and H<sub>2</sub>O. After extraction with ethylacetate (3 times), the combined organic phase was washed with brine, dried over MgSO<sub>4</sub> and concentrated. The residue was chromatographed on a column of silica gel eluting with hexane/ethyl acetate (2:1) to give 3.47 g (60%) of the title compound as a white solid.

MS (ESI) m/z: 195 (M+H<sup>+</sup>), 193 (M-H<sup>+</sup>).

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 7.06-6.99 (1 H, m), 6.90 (1 H, dd, J=9.2, 2.4 Hz), 6.82-6.72 (1 H,

m), 4.83-4.62 (1 H, m), 1.54 (6 H, d, J=7.1 Hz)

Step 3. 5-fluoro-N-([1-[(4-hydroxytetrahydro-2H-pyran-4-yl)methyl]piperidin-4-yl)methyl]-3-isopropyl-2-oxo-2,3-dihydro-1H-benzimidazole-1-carboxamide

To a stirred mixture of 6-fluoro-1-isopropyl-1,3-dihydro-2H-benzimidazol-2-one (Step 2 of Example 9, 0.58 g, 3 mmol) and p-nitrophenylchloroformate (0.66 g, 3.3 mmol) in dichloromethane (15 mL) was added triethylamine (1.25 mL, 9.0 mmol) at room temperature. After being stirred for 2h, a solution of 4-[[4-(aminomethyl)piperidin-1-yl]methyl]tetrahydro-2H-pyran-4-ol (Step 2 of Example 1, 0.75 g, 3.3 mmol) in dichloromethane (15 mL) was added to the mixture. After being stirred for 4h, the mixture was diluted with ethyl acetate (100 mL). Then, the organic layer was washed with 0.5 N NaOH aq. (10 mL) for 5 times and brine, dried over MgSO<sub>4</sub> and concentrated. The residue was chromatographed on a column of aminopropyl-silica gel eluting with hexane/ethyl acetate (3:1) to give 0.97 g (79 %) of the title compound as a white solid.


MS (ESI) m/z: 449 (M+H<sup>+</sup>).

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 8.84-8.74 (1 H, m), 8.21-8.11 (2 H, m), 7.02-6.91 (2 H, m), 4.68-4.56 (1 H, m), 3.87-3.72 (4 H, m), 3.34-3.25 (2 H, m), 2.93-2.82 (2 H, m), 2.42-2.25 (4 H, m), 1.79-1.68 (2 H, m), 1.67-1.29 (13 H, m).

Anal. calcd. for C<sub>23</sub>H<sub>33</sub>N<sub>4</sub>O<sub>4</sub>F: C, 61.59; H, 7.42; N, 12.49. Found: C, 61.45; H, 7.33; N, 12.40.

**EXAMPLE 10:**

5,6-difluoro-N-([1-[(4-hydroxytetrahydro-2H-pyran-4-yl)methyl]piperidin-4-yl)methyl]-3-isopropyl-2-oxo-2,3-dihydro-1H-benzimidazole-1-carboxamide hydrochloride



Step 1. 4,5-difluoro-N-isopropyl-2-nitroaniline

4,5-difluoro-2-nitroaniline (3.48 g, 20 mmol), 2,2-dimethoxypropane (11.9 mL, 100 mmol), and trifluoroacetic acid (1.6 mL, 21 mmol) were dissolved in toluene

(40 mL) and stirred at room temperature for 1 h. A boron-pyridine complex (2.12 mL, 21 mmol) was slowly added. The reaction mixture was stirred for 20 h. The solvent was evaporated in vacuo, and the residue was taken up into water and extracted with dichloromethane. The organic extract was dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated in vacuo.

- 5 The residue was chromatographed on a column of aminopropyl-silica gel eluting with hexane/ethyl acetate (30:1) to give 2.42 g (56%) of the title compound as a bright orange solid.

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 8.05 (1 H, dd, J=10.8, 8.6 Hz), 6.61 (1 H, dd, J=12.6, 6.8 Hz), 3.77-3.62 (1 H, m), 1.33 (6 H, d, J=6.2 Hz).

10 Step 2. 5,6-difluoro-1-isopropyl-1,3-dihydro-2H-benzimidazol-2-one

The title compound was prepared according to the procedure described in Step2 of Example 9 from 4,5-difluoro-*N*-isopropyl-2-nitroaniline (Step 1 of Example 10).

MS (ESI) m/z: 213 (M+H<sup>+</sup>), 211 (M+H<sup>+</sup>).

- 15 <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 7.00-6.89 (2 H, m), 4.76-4.57 (1 H, m), 3.86-3.69 (4 H, m), 3.31 (2 H, t, J=7.0 Hz), 2.95-2.82 (2 H, m), 2.35 (2 H, t, J=, 13.7 Hz), 2.31 (2 H, s), 1.67-1.25 (10 H, m), 1.55 (6 H, d, J=7.7 Hz).

Step 3. 5,6-difluoro-*N*-(1-[(4-hydroxytetrahydro-2H-pyran-4-yl)methyl]piperidin-4-yl)methyl)-3-isopropyl-2-oxo-2,3-dihydro-1H-benzimidazole-1-carboxamide

- 20 The title compound was prepared according to the procedure described in Step3 of Example 9 from 5,6-difluoro-1-isopropyl-1,3-dihydro-2H-benzimidazol-2-one (Step 2 of Example 10) and 4-[[4-(aminomethyl)piperidin-1-yl)methyl]tetrahydro-2H-pyran-4-ol(Step 2 of Example 1).

MS (ESI) m/z: 467 (M+H<sup>+</sup>).

- 25 <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 8.88-8.78 (1 H, m), 8.25-8.15 (1 H, m), 6.94-6.79 (2 H, m), 4.73-4.57 (1 H, m), 3.86-3.69 (4 H, m), 3.31 (2 H, t, J=7.0 Hz), 2.95-2.82 (2 H, m), 2.35 (2 H, t, J=, 13.7 Hz), 2.31 (2 H, s), 1.67-1.25 (10 H, m), 1.55 (6 H, d, J=7.7 Hz).

Step 4. 5,6-difluoro-*N*-(1-[(4-hydroxytetrahydro-2H-pyran-4-yl)methyl]piperidin-4-yl)methyl)-3-isopropyl-2-oxo-2,3-dihydro-1H-benzimidazole-1-carboxamide

- 30 hydrochloride

A mixture of 5,6-difluoro-*N*-(1-[(4-hydroxytetrahydro-2H-pyran-4-yl)methyl]piperidin-4-yl)methyl)-3-isopropyl-2-oxo-2,3-dihydro-1H-benzimidazole-

1-carboxamide (Step 3 of Example 10, 113 mg, 0.242 mmol) and 10% HCl-methanol (5 mL) was stirred for 1h. Then, the volatile components were removed under reduced pressure and the residue was recrystallized from ethanol-diethyl ether to give 88 mg (72 %) of the title compound as a colorless powder.

- 5 MS (ESI) m/z: 467 (M+H<sup>+</sup>).


<sup>1</sup>H NMR (DMSO-d<sub>6</sub>): δ 8.82-8.71 (1 H, m), 8.08-7.93 (1 H, m), 7.78-7.67 (1 H, m), 5.35-5.26 (1 H, m), 4.69-4.52 (1 H, m), 3.70-3.51 (6 H, m), 3.41-2.91 (7 H, m), 1.94-1.53 (8 H, m), 1.45 (6 H, d, J=7.0 Hz).

Anal. calcd. for C<sub>23</sub>H<sub>33</sub>N<sub>4</sub>O<sub>4</sub>F<sub>2</sub>Cl·1H<sub>2</sub>O: C, 53.96; H, 6.69; N, 10.94. Found: C,

- 10 53.67; H, 6.64; N, 10.89.

### **EXAMPLE 11:**

**6-chloro-N-({1-[4-hydroxytetrahydro-2H-pyran-4-yl)methyl]piperidin-4-yl)methyl}-3-isopropyl-2-oxo-2,3-dihydro-1H-benzimidazole-1-carboxamide hydrochloride**



**Step 1. 6-chloro-N-({1-[4-hydroxytetrahydro-2H-pyran-4-yl)methyl]piperidin-4-yl)methyl}-3-isopropyl-2-oxo-2,3-dihydro-1H-benzimidazole-1-carboxamide**

- 20 The title compound was prepared according to the procedure described in Step 3 of Example 9 from 5-chloro-1-isopropyl-1,3-dihydro-2H-benzimidazol-2-one (I. Tapia et al., *J. Med. Chem.*, 42, 2880 (1999)) and 4-{{[4-(aminomethyl)piperidin-1-yl)methyl]tetrahydro-2H-pyran-4-ol}(Step 2 of Example 1).

MS (ESI) m/z: 465 (M+H<sup>+</sup>).

- <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 8.33-8.30 (1 H, m), 7.19-7.14 (1 H, m), 7.04-7.03 (1 H, m), 4.73-4.57 (1 H, m), 3.82-3.71 (4 H, m), 3.31 (2 H, t, J=6.4 Hz), 2.95-2.83 (2 H, m), 2.41-2.29 (4 H, m), 1.79-1.68 (2 H, m), 1.67-1.25 (8 H, m), 1.54 (6 H, d, J=7.0 Hz).

**Step 2. 6-chloro-N-({1-[4-hydroxytetrahydro-2H-pyran-4-yl)methyl]piperidin-4-yl)methyl}-3-isopropyl-2-oxo-2,3-dihydro-1H-benzimidazole-1-carboxamide hydrochloride**



The title compound was prepared according to the procedure described in Step 4 of Example 10 from 6-chloro-*N*-({1-[(4-hydroxytetrahydro-2*H*-pyran-4-yl)methyl]piperidin-4-yl)methyl}-3-isopropyl-2-oxo-2,3-dihydro-1*H*-benzimidazole-1-carboxamide (Step 1 of Example 11)


5 MS (ESI) *m/z*: 465 (*M*+*H*<sup>+</sup>).

<sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>): δ 8.84-8.76 (1 H, m), 8.10-8.07 (1 H, m), 7.51-7.45 (1 H, m), 7.32-7.25 (1 H, m), 5.38-5.32 (1 H, m), 4.73-4.56 (1 H, m), 3.70-3.55 (6 H, m), 3.41-2.91 (7 H, m), 1.95-1.58 (8 H, m), 1.48 (6 H, d, *J*=7.7 Hz).

Anal. calcd. for C<sub>23</sub>H<sub>34</sub>N<sub>4</sub>O<sub>4</sub>Cl<sub>2</sub>·0.5H<sub>2</sub>O: C, 54.12; H, 6.91; N, 10.98. Found: C, 53.85; H, 6.90; N, 10.78.

#### EXAMPLE 12:

##### 5-chloro-*N*-({1-[(4-hydroxytetrahydro-2*H*-pyran-4-yl)methyl]piperidin-4-yl)methyl}-3-isopropyl-2-oxo-2,3-dihydro-1*H*-benzimidazole-1-carboxamide



##### Step 1. 5-chloro-*N*-isopropyl-2-nitroaniline

The title compound was prepared according to the procedure described in Step 1 of Example 10 from 5-chloro-2-nitroaniline.

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 8.12 (1 H, d, *J*=9.2 Hz), 6.84 (1 H, d, *J*=2.0 Hz), 6.57 (1 H, dd, *J*=9.2, 2.0 Hz), 3.81-3.71 (1 H, m), 1.33 (6 H, d, *J*=6.2 Hz)

##### Step 2. 6-chloro-1-isopropyl-1,3-dihydro-2*H*-benzimidazol-2-one

A mixture of 5-chloro-*N*-isopropyl-2-nitroaniline (Step 1 of Example 12, 0.76 g, 3.54 mmol), iron (0.99 g, 17.7 mmol) and ammonium chloride (0.38 g, 7.08 mmol) was suspended in ethanol (27 mL) and H<sub>2</sub>O (9 mL). Then, the mixture was heated at 80°C for 3h. After cooling, the insoluble materials was filtered off on a pad of Celite, and the filtrate was evaporated under reduced pressure. To the residue was added *N,N'*-carbonyldiimidazole (CDI, 0.57 g, 3.50 mmol) and THF (10 mL) and then stirred at 100°C for 10h. After cooling, the volatile materials were removed under reduced pressure and the residue was partitioned between ethylacetate and H<sub>2</sub>O. After

extraction with ethylacetate (3 times), the combined organic phase was washed with brine, dried over  $\text{MgSO}_4$  and concentrated. The residue was chromatographed on a column of silica gel eluting with hexane/ethyl acetate (2:1) to give 0.30 g (40%) of the title compound as a white solid..

- 5  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  6.99-6.90 (2 H, m), 6.84-6.74 (1 H, m), 4.94-4.77 (1 H, m), 1.64 (6 H, d,  $J=7.0$  Hz)

Step 3. 5-chloro-*N*-({1-[(4-hydroxytetrahydro-2*H*-pyran-4-yl)methyl]piperidin-4-yl)methyl}-3-isopropyl-2-oxo-2,3-dihydro-1*H*-benzimidazole-1-carboxamide

- The title compound was prepared according to the procedure described in Step 3 of Example 9 from 6-chloro-1-isopropyl-1,3-dihydro-2*H*-benzimidazol-2-one (Step 2 of Example 12) and 4-{{4-(aminomethyl)piperidin-1-yl}methyl}tetrahydro-2*H*-pyran-4-ol (Step 2 of Example 1).


MS (ESI)  $m/z$ : 465 ( $\text{M}+\text{H}^+$ ).

- 15  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  8.88-8.78 (1 H, m), 8.21-8.14 (1 H, m), 7.19-7.10 (2 H, m), 4.73-4.56 (1 H, m), 3.87-3.69 (4 H, m), 3.30 (2 H, t,  $J=6.2$  Hz), 2.94-2.84 (2 H, m), 2.41-2.27 (4 H, m), 1.79-1.68 (2 H, m), 1.67-1.25 (11 H, m).

Anal. calcd. for  $\text{C}_{23}\text{H}_{33}\text{N}_4\text{O}_4\text{Cl}$ : C, 59.41; H, 7.15; N, 12.05. Found: C, 59.27; H, 7.10; N, 11.72.

20 **EXAMPLE 13:**

*N*-({1-[(4-hydroxytetrahydro-2*H*-pyran-4-yl)methyl]piperidin-4-yl)methyl}-3-isopropyl-5-methyl-2-oxo-2,3-dihydro-1*H*-benzimidazole-1-carboxamide



Step 1. *N*-isopropyl-5-methyl-2-nitroaniline

- 25 The title compound was prepared according to the procedure described in Step 1 of Example 9 from 2-fluoro-4-methyl-1-nitrobenzene.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  8.12-8.01 (2 H, m), 6.63 (1 H, brs), 6.42 (1 H, d,  $J=10.3$  Hz), 3.94-3.72 (1 H, m), 2.33 (3 H, s), 1.32 (6 H, d,  $J=6.4$  Hz)

Step 2. 1-isopropyl-6-methyl-1,3-dihydro-2*H*-benzimidazol-2-one

The title compound was prepared according to the procedure described in Step2 of Example 9 from *N*-isopropyl-5-methyl-2-nitroaniline (Step 1 of Example 13). MS (ESI) *m/z*: 191 (*M*+*H*<sup>+</sup>).

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 7.04-6.93 (2 H, m), 6.90-6.80 (1 H, m), 4.82-4.63 (1 H, m), 2.40 (3 H, s), 1.55 (6 H, d, *J*=7.0 Hz).

Step 3. *N*-({1-[(4-hydroxytetrahydro-2*H*-pyran-4-yl)methyl]piperidin-4-yl)methyl}-3-isopropyl-5-methyl-2-oxo-2,3-dihydro-1*H*-benzimidazole-1-carboxamide

The title compound was prepared according to the procedure described in Step 3 of Example 9 from 1-isopropyl-6-methyl-1,3-dihydro-2*H*-benzimidazol-2-one (Step 2 of Example 13) and 4-{{4-(aminomethyl)piperidin-1-yl)methyl} tetrahydro-2*H*-pyran-4-ol(Step 2 of Example 1).


MS (ESI) *m/z*: 445 (*M*+*H*<sup>+</sup>).

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 8.97-8.84 (1 H, m), 8.10 (1 H, d, *J*= 8.8Hz), 7.01-6.93 (2 H, m), 4.76-4.58 (1 H, m), 3.85-3.69 (4 H, m), 3.30 (2 H, t, *J*=6.4 Hz), 2.94-2.82 (2 H, m), 2.41 (3 H, s), 2.43-2.27 (4 H, m), 1.80-1.68 (2 H, m), 1.67-1.25 (11 H, m).

Anal. calcd. for C<sub>24</sub>H<sub>36</sub>N<sub>4</sub>O<sub>4</sub>: C, 64.84; H, 8.16; N, 12.60. Found: C, 64.78; H, 8.29; N, 12.58.

#### EXAMPLE 14:

*N*-({1-[(4-hydroxytetrahydro-2*H*-pyran-4-yl)methyl]piperidin-4-yl)methyl}-3-isopropyl-4-methyl-2-oxo-2,3-dihydro-1*H*-benzimidazole-1-carboxamide



Step 1. *N*-({1-[(4-hydroxytetrahydro-2*H*-pyran-4-yl)methyl]piperidin-4-yl)methyl}-3-isopropyl-4-methyl-2-oxo-2,3-dihydro-1*H*-benzimidazole-1-carboxamide

The title compound was prepared according to the procedure described in Step 3 of Example 9 from 1-isopropyl-7-methyl-1,3-dihydro-2*H*-benzimidazol-2-one (I. Tapia et al., *J. Med. Chem.*, 42, 2880 (1999)) and 4-{{4-(aminomethyl)piperidin-1-yl)methyl} tetrahydro-2*H*-pyran-4-ol(Step 2 of Example 1).

MS (ESI) *m/z*: 445 (*M*+*H*<sup>+</sup>).

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 9.11-8.97 (1 H, m), 8.17 (1 H, d, J= 7.7Hz), 7.10-6.88 (2 H, m), 4.99-4.82 (1 H, m), 3.91-3.69 (4 H, m), 3.29 (2 H, t, J=6.2 Hz), 2.94-2.82 (2 H, m), 2.59 (3 H, s), 2.43-2.27 (4 H, m), 1.84-1.19 (7 H, m), 1.62 (6 H, d, J=6.8 Hz).


Anal. calcd. for C<sub>24</sub>H<sub>36</sub>N<sub>4</sub>O<sub>4</sub>: C, 64.84; H, 8.16; N, 12.60. Found: C, 64.73; H, 8.35;

5 N, 12.56.

### EXAMPLE 15:

*N-({1-[4-(4-hydroxytetrahydro-2H-pyran-4-yl)methyl]piperidin-4-yl}methyl)-3-isopropyl-4,5-dimethyl-2-oxo-2,3-dihydro-1H-benzimidazole-1-carboxamide*

#### 10 hydrochloride



#### Step 1. *N-isopropyl-2,3-dimethyl-6-nitroaniline*

The title compound was prepared according to the procedure described in Step 1 of Example 10 from 2,3-dimethyl-6-nitroaniline.

15 <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 7.82 (1 H, d, J=8.6 Hz), 6.79 (1 H, d, J=8.4 Hz), 3.52-3.34 (1 H, m), 2.30 (3 H, s), 2.24 (3 H, s), 1.11 (6 H, d, J=6.2 Hz)

#### Step 2. *1-isopropyl-6,7-dimethyl-1,3-dihydro-2H-benzimidazol-2-one*

The title compound was prepared according to the procedure described in Step2 of Example 9 from *N-isopropyl-2,3-dimethyl-6-nitroaniline* (Step 1 of Example 15).

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 7.11 (1 H, brs), 6.92-6.70 (1 H, m), 5.00-4.82 (1 H, m), 2.45 (3 H, s), 2.32 (3 H, s), 1.63 (6 H, d, J=7.0 Hz).

*Step 3. N-({1-[4-(4-hydroxytetrahydro-2H-pyran-4-yl)methyl]piperidin-4-yl}methyl)-3-isopropyl-4,5-dimethyl-2-oxo-2,3-dihydro-1H-benzimidazole-1-carboxamide*

#### 25 hydrochloride

To a stirred mixture of 1-isopropyl-6,7-dimethyl-1,3-dihydro-2H-benzimidazol-2-one (Step 2 of Example 15, 204 mg, 1 mmol) and p-nitrophenylchloroformate (220 mg, 1.1 mmol) in dichloromethane (7 mL) was added triethylamine (0.42 mL, 3.0 mmol) at room temperature. After being stirred for 2h, a

solution of 4-[[4-(aminomethyl)piperidin-1-yl]methyl]tetrahydro-2H-pyran-4-ol (Step 2 of Example 1, 230 mg, 1.0 mmol) in dichloromethane (3 mL) was added to the mixture. After being stirred for 4h, the mixture was diluted with ethyl acetate (50 mL). Then, the organic layer was washed with 0.5 N NaOH aq. (5 mL) for 5 times and brine, dried over MgSO<sub>4</sub> and concentrated. The residue was filtered through pad of aminopropyl-silica gel eluting with hexane/ethyl acetate (3:1) and the filtrate was concentrated. To the mixture was added 10% HCl-methanol (5 mL) was stirred for 1h. Then, the volatile components were removed under reduced pressure and the residue was recrystallized from ethanol-diethyl ether to give 100 mg (20 %) of the title compound as a colorless powder.


MS (ESI) m/z: 459 (M+H<sup>+</sup>).

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>): δ 8.96-8.87 (1 H, m), 7.84 (1 H, d, J=8.3 Hz), 6.95 (1 H, d, J=8.3 Hz), 5.34-5.21 (1 H, m), 5.01-4.86 (1 H, m), 3.69-3.53 (6 H, m), 3.41-2.91 (7 H, m), 2.45 (3 H, s), 2.28 (3 H, s), 1.87-1.70 (3 H, m), 1.67-1.48 (5 H, m), 1.52 (6 H, d, J=6.6 Hz).

Anal. calcd. for C<sub>25</sub>H<sub>39</sub>N<sub>4</sub>O<sub>4</sub>Cl·0.5H<sub>2</sub>O: C, 59.57; H, 8.00; N, 11.12. Found: C, 59.53; H, 7.98; N, 11.10.

#### **EXAMPLE 16:**

**6-fluoro-N-({1-[(4-hydroxytetrahydro-2H-pyran-4-yl)methyl]piperidin-4-yl)methyl}-3-isopropyl-5-methyl-2-oxo-2,3-dihydro-1H-benzimidazole-1-carboxamide hydrochloride**



#### **Step 1. 4-fluoro-N-isopropyl-5-methyl-2-nitroaniline**

The title compound was prepared according to the procedure described in Step 1 of Example 9 from 1,4-difluoro-2-methyl-5-nitrobenzene (T. Timothy et al., *J. Med. Chem.*, 35, 2321 (1992)).

MS (ESI) m/z: 213 (M+H<sup>+</sup>).

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 7.82 (1 H, d, J=10.3 Hz), 6.64 (1 H, d, J=6.4 Hz), 3.88-3.67 (1 H,

m), 2.30 (3 H, s), 1.31 (6 H, d, J=6.4 Hz)

Step 2. 5-fluoro-1-isopropyl-6-methyl-1,3-dihydro-2H-benzimidazol-2-one

The title compound was prepared according to the procedure described in Step2 of Example 9 from 4-fluoro-*N*-isopropyl-5-methyl-2-nitroaniline (Step 1 of Example 16).

MS (ESI) m/z: 209 (M+H<sup>+</sup>).

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 7.00-6.96 (1 H, m), 6.92-6.90 (1 H, m), 4.75-4.56 (1 H, m), 2.31 (3 H, s), 1.55 (6 H, d, J=7.0 Hz).

Step 3. 6-fluoro-*N*-({1-[(4-hydroxytetrahydro-2H-pyran-4-yl)methyl]piperidin-4-yl)methyl}-3-isopropyl-5-methyl-2-oxo-2,3-dihydro-1H-benzimidazole-1-carboxamide

The title compound was prepared according to the procedure described in Step 3 of Example 9 from 5-fluoro-1-isopropyl-6-methyl-1,3-dihydro-2H-benzimidazol-2-one (Step 2 of Example 16) and 4-{{[4-(aminomethyl)piperidin-1-yl]methyl}tetrahydro-2H-pyran-4-ol} (Step 2 of Example 1).

MS (ESI) m/z: 463 (M+H<sup>+</sup>).

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 8.92-8.83 (1 H, m), 7.96 (1 H, d, J=10.1 Hz), 6.91 (1 H, d, J=6.2 Hz), 4.75-4.56 (1 H, m), 3.85-3.70 (4 H, m), 3.30 (2 H, t, J=6.4 Hz), 2.94-2.82 (2 H, m), 2.42-2.29 (7 H, m), 1.84-1.19 (7 H, m), 1.55 (6 H, d, J=7.0 Hz).

Step 4. 6-fluoro-*N*-({1-[(4-hydroxytetrahydro-2H-pyran-4-yl)methyl]piperidin-4-yl)methyl}-3-isopropyl-5-methyl-2-oxo-2,3-dihydro-1H-benzimidazole-1-carboxamide hydrochloride

The title compound was prepared according to the procedure described in Step 4 of Example 10 from 6-fluoro-*N*-({1-[(4-hydroxytetrahydro-2H-pyran-4-yl)methyl]piperidin-4-yl)methyl}-3-isopropyl-5-methyl-2-oxo-2,3-dihydro-1H-benzimidazole-1-carboxamide (Step 3 of Example 16)

MS (ESI) m/z: 463 (M+H<sup>+</sup>).

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>): δ 9.55-9.11 (1 H, m), 8.89-8.74 (1 H, m), 7.77 (1 H, d, J=10.4 Hz), 7.40 (1 H, d, J=6.6 Hz), 5.42-5.34 (1 H, m), 4.70-4.56 (1 H, m), 3.69-3.53 (6 H, m), 3.52-2.91 (7 H, m), 2.29 (3 H, s), 1.87-1.70 (3 H, m), 1.95-1.55 (8 H, m), 1.48 (6 H, d, J=6.8 Hz).


Anal. calcd. for C<sub>24</sub>H<sub>36</sub>N<sub>4</sub>O<sub>4</sub>·HCl: C, 57.76; H, 7.27; N, 11.23. Found: C, 57.47; H,

7.40; N, 11.05.

### PREPARATION 1.

Step 1. *tert*-Butyl 4-(((3-isopropyl-2-oxo-2,3-dihydro-1*H*-benzimidazol-1-

5 yl)carbonyl)amino)methyl)piperidine-1-carboxylate




To a stirred solution of 1-isopropyl-1,3-dihydro-2*H*-benzimidazol-2-one (*J. Med. Chem.* **1999**, *42*, 2870-2880) (3.00 g, 17.02 mmol) and triethylamine (7.12 ml, 51.06 mmol) in 70 ml tetrahydrofuran was added triphosgen (5.15 g, 17.02 mmol) in 14 ml tetrahydrofuran at room temperature. The reaction mixture was refluxed for 19

10 hours. The mixture was then cooled to room temperature, *tert*-butyl 4-(aminomethyl)piperidine-1-carboxylate (*J. Prugh, L. A. Birchenough and M. S. Egberton, Synth. Commun.*, **1992**, *22*, 2357-60) (3.28 g, 15.32 mmol) in 10 ml tetrahydrofuran was added. The reaction mixture was refluxed for another 24 hours. Then cooled and basified with aqueous saturated NaHCO<sub>3</sub> 50 ml, and extracted with ethyl acetate 100 ml for three times. The combined extract was washed with brine,

15 dried over MgSO<sub>4</sub> and concentrated. Flash chromatography of the residue (eluent: hexane/ethyl acetate=5/1 to 1/2) afforded a colorless oil 3.99 g (62%) as the titled compound.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 9.04-8.88 (1 H, m), 8.83-8.20 (1H, m), 7.26-7.10 (3H, m), 4.80-4.60 (1H, m), 4.28-4.02 (2H, m), 3.32 (2H, t, J=6.1 Hz), 2.82-2.60 (2H, m), 1.94-1.10 (5H, m), 1.57 (6H, d, J=7.1 Hz), 1.45 (9H, s).

Step 2. 3-Isopropyl-2-oxo-*N*-(piperidin-4-ylmethyl)-2,3-dihydro-1*H*-benzimidazole-1-carboxamide




A solution of tert-butyl 4-({[(3-isopropyl-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl)carbonyl]amino}methyl)piperidine-1-carboxylate (3.992 g, 9.58 mmol) in 50 ml 10% hydrochloric acid in methanol and 10 ml concentrated hydrochloric acid was stirred at room temperature for 18 hours. The mixture was then concentrated and basified with aqueous Na<sub>2</sub>CO<sub>3</sub>, extracted with CHCl<sub>3</sub> 100 ml for 3 times. The combined extract was dried and concentrated. Flash chromatography of the residue (NH-silica gel, eluent: CH<sub>2</sub>Cl<sub>2</sub>/methanol=100/1) afforded a colorless oil 2.272 g (75%) as the titled compound.

MS (ESI) m/z: 317 (M+H)<sup>+</sup>.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 8.93 (1H, br), 8.32-8.22 (1H, m), 7.24-7.02 (3H, m), 4.80-4.61 (1H, m), 3.31 (2H, t, J=6.0 Hz), 3.20-3.05 (2H, m), 2.79-2.54 (2H, m), 1.84-1.52 (3H, m), 1.57 (6H, d, J=6.9 Hz), 1.36-1.13 (2H, m).

Step 3. N-([1-(3-Hydroxy-3-methyl-2-oxobutyl)piperidin-4-yl]methyl)-3-isopropyl-2-oxo-2,3-dihydro-1H-benzimidazole-1-carboxamide, monooxalate salt



A mixture of 3-isopropyl-2-oxo-N-(piperidin-4-ylmethyl)-2,3-dihydro-1H-benzimidazole-1-carboxamide (250 mg, 0.790 mmol), 1-bromo-3-hydroxy-3-methylbutan-2-one (G. Bertram; A. Scherer; W. Steglich; W. Weber, *Tetrahedron Lett.*, 1996, 37, 7955-7958) (181 mg, 1.343 mmol) and triethylamine (0.28 ml, 1.975 mmol) in 8 ml tetrahydrofuran was refluxed for 15 hours. Then cooled and diluted with 100 ml ethyl acetate and was washed with aqueous NaHCO<sub>3</sub> 20 ml, brine, dried over MgSO<sub>4</sub> and concentrated. Flash chromatography of the residue (eluent: CH<sub>2</sub>Cl<sub>2</sub>/methanol=100/1 to 30/1) afforded a colorless oil 202 mg (61%). The oil (202 mg) was dissolved in 3 ml methanol and acidified with a solution of 44 mg oxalic acid in 1 ml MeOH. The mixture was concentrated. Recrystallization of the resulting solid with EtOH-AcOEt afforded a white solid 246 mg as the titled compound.

MS (ESI) m/z: 417 (M+H)<sup>+</sup>.

m.p.: 140.5°C.



IR (KBr)  $\nu$ : 3404, 3306, 2980, 2941, 1728, 1690, 1612, 1541  $\text{cm}^{-1}$ .


$^1\text{H-NMR}$  ( $\text{CDCl}_3$ ) (free base)  $\delta$ : 8.90 (1H, br) 8.30-8.20 (1H, m), 7.24-7.10 (3H, m), 4.78-4.61 (1H, m), 3.37 (2H, s), 3.33 (2H, t,  $J=6.3$  Hz), 3.00-2.86 (2H, m), 2.22-2.06 (2H, m), 1.90-1.22 (5H, m), 1.57 (6H, d,  $J=7.0$  Hz), 1.35 (6H, s).

- 5  $^1\text{H-NMR}$  ( $\text{DMSO-d}_6$ ) (salt form)  $\delta$ : 8.92-8.81 (1H, m) 8.07 (1H, dd,  $J=7.7, 6.8$  Hz), 7.44 (1H, d,  $J=7.7$  Hz), 7.28-7.10 (2H, m), 4.74-4.60 (1H, m), 4.36 (2H, bv), 4.00-2.70 (6H, m), 1.90-1.44 (5H, m), 1.49 (6H, d,  $J=6.4$  Hz), 1.24 (6H, s).

Anal. Calcd. for  $\text{C}_{24}\text{H}_{34}\text{N}_4\text{O}_8 \cdot 0.3\text{C}_2\text{H}_6\text{O} \cdot \text{H}_2\text{O}$ : C, 54.88; H, 7.08; N, 10.41. Found: C, 55.26; H, 7.18; N, 10.07.

## CLAIMS

1. A compound of the formula (I):



wherein


- 5 **Het** represents a heterocyclic group having one nitrogen atom, to which B binds directly, and from 4 to 7 carbon atoms, and said heterocyclic group being unsubstituted or substituted by 1 to 4 substituents independently selected from the group consisting of substituents  $\alpha^1$ ;
- A** represents an alkylene group having from 1 to 4 carbon atoms;
- 10 **B** represents a covalent bond or an alkylene group having from 1 to 5 carbon atoms, and said alkylene group being unsubstituted or substituted by an oxo group when  $R^3$  represents a heterocyclic group;
- R<sup>1</sup>** represents an isopropyl group or a cyclopentyl group;
- R<sup>2</sup>** independently represents a halogen atom or an alkyl group having from 1 to 4
- 15 carbon atoms; **m** is 0, 1, 2, 3 or 4; and
- R<sup>3</sup>** represents
- (i) a cycloalkyl group having from 3 to 8 carbon atoms, and said cycloalkyl group being substituted by 1 to 5 substituents independently selected from the group consisting of substituents  $\alpha^2$ , or
  - 20 (ii) a heterocyclic group having from 3 to 8 atoms, and said heterocyclic group being unsubstituted or substituted by 1 to 5 substituents independently selected from the group consisting of substituents  $\beta$ ,
- said substituents  $\alpha^1$**  are independently selected from a hydroxy group and an amino group;
- 25 **said substituents  $\alpha^2$**  are independently selected from a hydroxy group, an amino group, a hydroxy-substituted alkyl group having from 1 to 4 carbon atoms, a carboxyl

group and an alkoxy group having from 1 to 4 carbon atoms; and

said substituents  $\beta$  are independently selected from a hydroxy group, a hydroxy-substituted alkyl group having from 1 to 4 carbon atoms, a carboxyl group, an amino group, an alkyl group having from 1 to 4 carbon atoms, an amino-substituted alkyl group having from 1 to 4 carbon atoms and a carbamoyl group, or a pharmaceutically acceptable salts thereof.

2. The compound or the pharmaceutically acceptable salt thereof, as claimed in claim 1, wherein


10 **Het** represents a heterocyclic group selected from



said heterocyclic group being unsubstituted or substituted by 1 to 3 substituents independently selected from the group consisting of substituents  $\alpha^1$ ; and

**A** represents an alkylene group having from 1 to 3 carbon atoms.

3. The compound or the pharmaceutically acceptable salt thereof, as claimed in claim 1, wherein



**Het** represents a group of formula

and this group being unsubstituted or substituted by one substituent selected from the group consisting of substituents  $\alpha^1$ ;

**A** represents an alkylene group having from 1 to 2 carbon atoms;


**B** represents an alkylene group having from 1 to 4 carbon atoms, and said alkylene group being unsubstituted or substituted by an oxo group when  $R^3$  represents a heterocyclic group;

$R^2$  independently represents a halogen atom or an alkyl group having from 1 to 2 carbon atoms; **m** is 0, 1 or 2; and

**R<sup>3</sup>** represents

- (i) a cycloalkyl group having from 4 to 7 carbon atoms, and said cycloalkyl group being substituted by 1 to 3 substituents independently selected from the group consisting of substituents  $\alpha^2$ , or
- 5 (ii) a heterocyclic group having from 4 to 7 atoms, and said heterocyclic group being unsubstituted or substituted by 1 to 3 substituents independently selected from the group consisting of substituents  $\beta$ .

4. The compound or the pharmaceutically acceptable salt thereof, as claimed in claim 1, wherein
- 10



**Het** represents a group of formula

and this group being unsubstituted or substituted by one substituent selected from the group consisting of substituents  $\alpha^1$ ;

**A** represents a methylene group;

- 15 **B** represents an alkylene group having from 1 to 2 carbon atoms;

**R<sup>1</sup>** represents an isopropyl group;

**R<sup>2</sup>** independently represents a fluorine atom, a chlorine atom or a methyl; and


**R<sup>3</sup>** represents

- (i) a cycloalkyl group having from 5 to 7 carbon atoms, and said cycloalkyl group being substituted by 1 to 2 substituents independently selected from the group consisting of substituents  $\alpha^2$ , or
- 20 (ii) a heterocyclic group having from 5 to 7 atoms, and said heterocyclic group being unsubstituted or substituted by 1 to 2 substituents independently selected from the group consisting of substituents  $\beta$ ,

- 25 **said substituents  $\alpha^2$**  are independently selected from a hydroxy group, an amino group and an alkoxy group having from 1 to 2 carbon atoms; and

- said substituents  $\beta$**  are independently selected from a hydroxy group, a hydroxy-substituted alkyl group having from 1 to 2 carbon atoms, a carboxyl group, an amino group, an amino-substituted alkyl group having from 1 to 2 carbon atoms and a
- 30 carbamoyl group.

5. The compound or the pharmaceutically acceptable salt thereof, as claimed in claim 1, wherein



Het represents a group of formula

5 A represents a methylene group;

B represents a methylene group;

R<sup>1</sup> represents an isopropyl group;

R<sup>2</sup> represents a fluorine atom; m is 0 or 1; and

R<sup>3</sup> represents

10 (i) a cycloalkyl group having from 5 to 6 carbon atoms, and said cycloalkyl group being substituted by 1 to 2 substituents independently selected from the group consisting of substituents  $\alpha^2$ , or

(ii) a heterocyclic group having from 5 to 6 atoms, and said heterocyclic group being unsubstituted or substituted by 1 to 2 substituents independently


15 selected from the group consisting of substituents  $\beta$ ,

said substituents  $\alpha^2$  are independently selected from a hydroxy group and an amino group; and

said substituents  $\beta$  are independently selected from a hydroxy group and an amino group.

20

6. The compound or the pharmaceutically acceptable salt thereof, as claimed in claim 1, wherein



Het represents a group of formula

A represents a methylene group;

25 B represents a methylene group;

R<sup>1</sup> represents an isopropyl group;

R<sup>2</sup> represents a fluorine atom; m is 0; and

R<sup>3</sup> represents

(i) a cyclohexyl group substituted by 1 to 2 substituents independently

selected from a hydroxy group or an amino group, or

- (ii) a heterocyclic group having from 6 atoms, and said heterocyclic group being substituted by a hydroxy group or an amino group.

- 5 7. The compound or the pharmaceutically acceptable salt, as claimed in claim 6, wherein

$R^3$  represents

- (i) a cyclohexyl group substituted by 1 or 2 hydroxy group, or  
(ii) a tetrahydropyran group substituted by 1 or 2 hydroxy group.

10

8. The compound or the pharmaceutically acceptable salt, as claimed in claim 7, wherein

$R^3$  represents hydroxytetrahydropyranyl or dihydroxycyclohexyl.

- 15 9. The compound as claimed in claim 1, which is selected from:

*N*-({1-[(4-hydroxytetrahydro-2*H*-pyran-4-yl)methyl]piperidin-4-yl)methyl}-3-isopropyl-2-oxo-2,3-dihydro-1*H*-benzimidazole-1-carboxamide;

*N*-({1-[(*trans*-1,4-dihydroxyhexyl)methyl]piperidin-4-yl)methyl}-3-isopropyl-2-oxo-2,3-dihydro-1*H*-benzimidazole-1-carboxamide;

- 20 *N*-({1-[(*cis*-1,4-dihydroxyhexyl)methyl]piperidin-4-yl)methyl}-3-isopropyl-2-oxo-2,3-dihydro-1*H*-benzimidazole-1-carboxamide; and  
6-fluoro-*N*-({1-[(4-hydroxytetrahydro-2*H*-pyran-4-yl)methyl]piperidin-4-yl)methyl}-3-isopropyl-2-oxo-2,3-dihydro-1*H*-benzimidazole-1-carboxamide,  
or the pharmaceutically acceptable salt thereof.

25

10. The compound or the pharmaceutically acceptable salt, as claimed in any one of claims 1 to 9, wherein the pharmaceutically acceptable salt is a hydrochloride or hemiedysilate.

- 30 11. A pharmaceutical composition including the compound or the pharmaceutically acceptable salt thereof, as claimed in any one of claims 1 to 10, together with a pharmaceutically acceptable excipient.

12. A method of the treatment of disease conditions mediated by 5-HT<sub>4</sub> receptor activity, in a mammalian subject, which comprises administering to the said subject in need of such treatment a therapeutically effective amount of the compound or the  
5 pharmaceutically acceptable salt thereof, as claimed in any one of claims 1 to 10.

13. A method of the treatment of diseases selected from gastroesophageal reflux disease, gastrointestinal disease, gastric motility disorder, non-ulcer dyspepsia, functional dyspepsia, irritable bowel syndrome (IBS), constipation, dyspepsia,  
10 esophagitis, gastroesophageal disease, nausea, central nervous system disease, Alzheimer's disease, cognitive disorder, emesis, migraine, neurological disease, pain, cardiovascular disorders, cardiac failure, heart arrhythmia, diabetes and apnea syndrome, which comprises administering to the said subject a therapeutically effective amount of the compound or the pharmaceutically acceptable salt thereof, as  
15 claimed in any one of claims 1 to 10.


14. Use of the compound or the pharmaceutically acceptable salt thereof, as claimed in any one of claims 1 to 10 for the manufacture of a medicament for the treatment of disease conditions mediated by 5-HT<sub>4</sub> receptor activity, in a mammalian subject.

20

15. Use of a compound as claimed in claim 14, wherein said condition is selected from gastroesophageal reflux disease, gastrointestinal disease, gastric motility disorder, non-ulcer dyspepsia, functional dyspepsia, irritable bowel syndrome (IBS), constipation, dyspepsia, esophagitis, gastroesophageal disease, nausea, central  
25 nervous system disease, Alzheimer's disease, cognitive disorder, emesis, migraine, neurological disease, pain, cardiovascular disorders, cardiac failure, heart arrhythmia, diabetes and apnea syndrome.

16. A compound of the formula (2-A'):

30



wherein

**R<sup>a</sup>** represents a hydrogen atom or a N-protecting group;

**Het** represents a heterocyclic group having one nitrogen atom, to which B binds directly, and from 4 to 7 carbon atoms, and said heterocyclic group being  
 5 unsubstituted or substituted by 1 to 4 substituents independently selected from the group consisting of substituents  $\alpha^1$ ;

**A** represents an alkylene group having from 1 to 4 carbon atoms;

**B** represents a covalent bond or an alkylene group having from 1 to 5 carbon atoms,  
 10 and said alkylene group being unsubstituted or substituted by an oxo group when **R<sup>3</sup>** represents a heterocyclic group;

**R<sup>3</sup>** represents

- (i) a cycloalkyl group having from 3 to 8 carbon atoms, and said cycloalkyl group being substituted by 1 to 5 substituents independently selected from  
 15 the group consisting of substituents  $\alpha^2$ , or
- (ii) a heterocyclic group having from 3 to 8 atoms, and said heterocyclic group being unsubstituted or substituted by 1 to 5 substituents independently selected from the group consisting of substituents  $\beta$ ,


**said substituents  $\alpha^1$**  are independently selected from a hydroxy group and an amino  
 20 group;

**said substituents  $\alpha^2$**  are independently selected from a hydroxy group, an amino group, a hydroxy-substituted alkyl group having from 1 to 4 carbon atoms, a carboxyl group and an alkoxy group having from 1 to 4 carbon atoms; and

**said substituents  $\beta$**  are independently selected from a hydroxy group, a hydroxy-substituted alkyl group having from 1 to 4 carbon atoms, a carboxyl group, an amino  
 25 group, an alkyl group having from 1 to 4 carbon atoms, an amino-substituted alkyl group having from 1 to 4 carbon atoms and a carbamoyl group,  
 or a salt thereof



17. The compound or the salt thereof, as claimed in claim 16, wherein  $R^a$  represents a hydrogen atom or a *t*-butoxycarbonyl group;



Het represents a group of formula

A represents an methylene group; B represents an methylene group; and

5  $R^3$  represents hydroxytetrahydropyranyl or dihydroxycyclohexyl.

18. A combination of the compound or the pharmaceutically acceptable salt thereof as claimed in any one of claims 1 to 10, and another pharmacologically active agent.

10 19. A pharmaceutical composition including the compound or the pharmaceutically acceptable salt thereof as claimed in any one of claims 1 to 10, and another pharmacologically active agent.

## INTERNATIONAL SEARCH REPORT

International Application No  
PC17/IB2004/002741

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 7 C07D405/14 C07D413/14 C07D401/12 C07D405/06 A61K31/4184  
A61P1/04

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, BEILSTEIN Data, CHEM ABS Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with Indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 00/37461 A (BOSMANS JEAN PAUL R M ; CLEYN MICHEL ANNA JOZEF DE (BE); JANSSEN PHARM) 29 June 2000 (2000-06-29) abstract	1
X	page 26 - page 27; examples A.6,A.7 page 17; figure V	16
X	EP 0 522 914 A (SYNTHELABO) 13 January 1993 (1993-01-13) page 4, paragraphs page 7, line 54; example 4 page 8, line 40; example 5 claims	16
X	DE 33 36 024 A (BOEHRINGER INGELHEIM KG) 18 April 1985 (1985-04-18) page 23; example 11 claims	16
-/-		



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

## \* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*I\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*Z\* document member of the same patent family

Date of the actual completion of the international search

14 December 2004

Date of mailing of the international search report

04/01/2005

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2340, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Stix-Malaun, E

## INTERNATIONAL SEARCH REPORT

International Application No.

PCT/IB2004/002741

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DE 23 19 814 A (CIBA GEIGY AG) 31 October 1973 (1973-10-31) claims -----	16
X	WO 02/053568 A (NISHIDA HIDEMITSU ; MOCHIDA PHARM CO LTD (JP); MUKAIHIRA TAKAFUMI (JP)) 11 July 2002 (2002-07-11) RN 441790-70-1,4-Piperidinol,4-(aminomethyl)-1 -(4-pyridinyl) (CA INDEX NAME) page 239; compound II -----	16
Y	WO 93/18027 A (SMITHKLINE BEECHAM PLC) 16 September 1993 (1993-09-16) cited in the application claims page 11; examples e13-e15 -----	1-19
Y	J.MED.CHEM., vol. 42, no. 15, 1999, pages 2870-2880, XP002310533 abstract page 2875; table 3; compound 5K page 2872; figure 2 -----	1-19

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/IB2004/002741

## Box II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:  
  
Although claims 11,12 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 8.4(a).

## Box III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.  
☐ No protest accompanied the payment of additional search fees.

## INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No.

PCT/IB2004/002741

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 0037461	A	29-06-2000	AU 770397 B2 19-02-2004
			AU 2432800 A 12-07-2000
			BG 105571 A 31-01-2002
			BR 9916491 A 04-09-2001
			CA 2355857 A1 29-06-2000
			CN 1331690 T 16-01-2002
			CZ 20012117 A3 17-10-2001
			EA 3608 B1 26-06-2003
			EE 200100335 A 15-08-2002
			WO 0037461 A1 29-06-2000
			EP 1140915 A1 10-10-2001
			HR 20010445 A1 30-06-2002
			HU 0104838 A2 29-07-2002
			ID 28953 A 19-07-2001
			JP 2002533337 T 08-10-2002
			NO 20012858 A 08-06-2001
			NZ 512871 A 26-11-2002
			PL 348417 A1 20-05-2002
			SK 8592001 A3 05-02-2002
			TR 200101962 T2 22-04-2002
			TW 570920 B 11-01-2004
			US 2003181456 A1 25-09-2003
			US 6544997 B1 08-04-2003
			ZA 200105135 A 21-06-2002
EP 0522914	A	13-01-1993	FR 2678268 A1 31-12-1992
			FR 2691147 A1 19-11-1993
			AT 124942 T 15-07-1995
			AU 645697 B2 20-01-1994
			AU 1858692 A 14-01-1993
			AU 645429 B2 13-01-1994
			AU 1858792 A 07-01-1993
			CA 2072434 A1 28-01-1993
			CA 2072529 A1 28-01-1993
			CN 1067887 A 13-01-1993
			CN 1067888 A 13-01-1993
			CZ 9202000 A3 13-01-1993
			CZ 9202001 A3 13-01-1993
			DE 69203401 D1 17-08-1995
			DE 69203401 T2 07-03-1996
			DK 520882 T3 11-12-1995
			EP 0522914 A1 13-01-1993
			EP 0520882 A1 30-12-1992
			ES 2077376 T3 16-11-1995
			FI 922980 A 28-12-1992
			FI 922981 A 28-12-1992
			GR 3017612 T3 31-01-1996
			HU 61740 A2 01-03-1993
			HU 61745 A2 01-03-1993
			IE 922091 A1 30-12-1992
			IE 922092 A1 30-12-1992
			JP 7173152 A 11-07-1995
			JP 3159525 B2 23-04-2001
			JP 5194419 A 03-08-1993
			MX 9203332 A1 01-12-1992
			MX 9203453 A1 01-12-1992
			NO 922522 A 28-12-1992
			NO 922523 A 28-12-1992

## INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No.

PC17IB2004/002741

Patent document cited in search report		Publication date		Patent family member(s)		Publication date
EP 0522914	A		PL	295042 A1		08-03-1993
			PL	295046 A1		22-03-1993
			US	5246939 A		21-09-1993
			US	5254560 A		19-10-1993
			ZA	9204783 A		31-03-1993
			ZA	9204784 A		31-03-1993
DE 3336024	A	18-04-1985	DE	3336024 A1		18-04-1985
			AR	243503 A1		31-08-1993
			AT	42544 T		15-05-1989
			AU	565839 B2		01-10-1987
			AU	3383484 A		18-04-1985
			CA	1242193 A1		20-09-1988
			CS	8501537 A2		15-08-1985
			CS	9104108 A3		12-08-1992
			DD	227963 A5		02-10-1985
			DE	3477922 D1		01-06-1989
			DK	474684 A , B,		05-04-1985
			EP	0136658 A2		10-04-1985
			ES	8505652 A1		01-10-1985
			ES	8601877 A1		01-03-1986
			ES	8601878 A1		01-03-1986
			FI	843856 A , B,		05-04-1985
			GR	80546 A1		05-02-1985
			HK	75892 A		09-10-1992
			HU	35641 A2		29-07-1985
			IE	58155 B1		14-07-1993
			IL	73125 A		31-10-1988
			JP	2007402 C		11-01-1996
			JP	4261152 A		17-09-1992
			JP	7039387 B		01-05-1995
			JP	1722993 C		24-12-1992
			JP	4006684 B		06-02-1992
			JP	60094959 A		28-05-1985
			KR	9107882 B1		04-10-1991
			MX	9203259 A1		01-07-1992
			NO	843979 A , B,		09-04-1985
			NZ	209759 A		26-04-1989
			PH	23764 A		03-11-1989
			PL	249852 A1		10-09-1985
			PL	253280 A1		11-02-1986
			PT	79306 A , B		01-11-1984
			SG	44192 G		04-09-1992
			SU	1373318 A3		07-02-1988
			SU	1306472 A3		23-04-1987
			US	4833140 A		23-05-1989
			US	4996224 A		26-02-1991
			US	5073671 A		17-12-1991
			US	5149808 A		22-09-1992
			US	4767759 A		30-08-1988
			YU	169684 A1		31-10-1987
			YU	199086 A1		31-10-1987
			ZA	8407755 A		25-06-1986
DE 2319814	A	31-10-1973	CH	566987 A5		30-09-1975
			DE	2319814 A1		31-10-1973
			FR	2181058 A1		30-11-1973
			GB	1410533 A		15-10-1975

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/IB2004/002741

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
DE 2319814 A		JP 49014474 A	07-02-1974
		US 3945953 A	23-03-1976
WO 02053568 A	11-07-2002	CA 2433174 A1	11-07-2002
		EP 1346994 A1	24-09-2003
		WO 02053568 A1	11-07-2002
		US 2003045520 A1	06-03-2003
		US 2004063716 A1	01-04-2004
WO 9318027 A	16-09-1993	AU 666357 B2	08-02-1996
		AU 3571893 A	05-10-1993
		CA 2131381 A1	16-09-1993
		EP 0629198 A1	21-12-1994
		WO 9318027 A1	16-09-1993
		JP 7504428 T	18-05-1995
		MX 9301157 A1	29-07-1994
		NZ 249217 A	25-06-1996
		ZA 9301430 A	22-10-1993